arXiv:1210.1597v1 [math.QA] 4 Oct 2012

A GLOBAL QUANTUM DUALITY PRINCIPLE FOR SUBGROUPS AND HOMOGENEOUS SPACES

Nicola CICCOLI$^\flat$, Fabio GAVARINI$^\#$

$^\flat$ *Dipartimento di Matematica e Informatica, Università di Perugia*
via Vanvitelli 1 — I-06123 Perugia, Italy
e-mail: ciccoli@dipmat.unipg.it

$^\#$ *Dipartimento di Matematica, Università di Roma "Tor Vergata"*
via della ricerca scientifica 1 — I-00133 Roma, Italy
e-mail: gavarini@mat.uniroma2.it

Abstract

We develop a quantum duality principle for subgroups and homogeneous spaces of a Poisson group G and its Poisson dual G^* in the context of *global quantizations*.

Starting from a quantum subgroup, resp. a quantum homogeneous space (both suitably defined, even in a weak sense), for a given quantum group (on G), we provide functorial recipes which produce a quantization of a quantum homogeneous space, resp. a quantum subgroup, for G^*. The outcome objects are quantizations in a stronger sense, so that the homogeneous spaces, resp. the subgroups, which come out as their specializations actually are coisotropic, resp. Poisson quotients.

If we start with quantizations in the strongest possible sense — in more or less common terminology, they are called *flat* — the outcome is strong too. In this case, subgroups are necessarily coisotropic and homogeneous spaces are Poisson quotients; also, in our construction any final object is related to the initial one by being its *coisotropic dual*. We end with some explicit examples of application of our recipes. [1]

1 Introduction

The quantum duality principle, first mentioned by Drinfel'd in [6] (see [8] for a proof), was a remark that quantized universal enveloping algebra could be interpreted also as quantum function algebras of suitable dual groups, and viceversa.

This principle — QDP in what follows — was generalized to (embeddable) quantum homogeneous spaces in [4]; to be more precise, in that work it was shown that in the context of formal quantization one can define functors on two kind of algebraic subobjects of a Hopf algebra: one-sided coideal subalgebras and one-sided ideals and coideals. These algebraic objects, depending on whether they sit inside a quantized enveloping algebra or a quantized function algebra, describe quantizations

---

[1] Keywords: *Quantum Groups, Poisson Homogeneous Spaces, Coisotropic Subgroups.*
2000 *Mathematics Subject Classification:* Primary 17B37, 20G42, 58B32; Secondary 81R50.

of coisotropic subgroups or quantizations of quotient Poisson homogeneous spaces. Such subobjects have been the subject of intensive study in last years, leading to their classification in some specific cases (e.g. [12, 13, 15]). The above mentioned functors transform then a quantum homogeneous space (resp. a quantum coisotropic subgroup) in a quantum coisotropic subgroup (resp. a quantum homogeneous space) for the dual Poisson group. Of course this duality, much like the original Drinfel'd's QDP, is dictated by the underlying Poisson structures.

The results of the present paper extend those in [4, 5], which deals with formal quantum groups and flat (deformation) quantizations of their subgroups and homogeneous spaces. Besides adapting that work to the richer context of global quantum groups, we now deal with weakest notions of quantizations of subgroup and homogeneous spaces: this allows to consider also those subgroups which are not coisotropic and those homogeneous spaces which are not Poisson quotients.

We will show how the generalized QDP extends from formal to *global* quantizations, i.e. suitable Hopf algebras over $\mathbb{C}\big[q,q^{-1}\big]$ and subobjects inside them (to be specialized at $q = 1$). The global QDP will then appear in the form of four functors transforming one-sided coideal subalgebras (resp. one-sided ideals and coideals) in a quantized universal enveloping algebra (resp. in a quantum function algebra) into one-sided coideal subalgebras (resp. one-sided ideals and coideals) in a quantum function algebra (resp. in a quantum universal enveloping algebra). The intermediate tool is the global version of QDP for quantum groups, developed in [10].

The first point to be remarked is that, since we consider global geometrical features, we profit of all advantages of such a richer point of view. On the other hand, it may happen (already at the classical level) that the usual subgroup-homogeneous space correspondence is lost, due to a lack in globally defined algebraic functions.

Secondly, we elaborate on the notion of (global) *quantization* of subobjects. We consider three versions of it, from the strongest — usually referred to as "flat" — to the weakest one, and we show how the QDP functors behave for each of them. The main advantage is the following: while the two strongest notions only apply to subgroups, resp. homogeneous space, which are *coisotropic*, resp. are *Poisson quotients*, the weakest one applies to any possible subgroup or homogeneous space. This definitely extends the range of application of the QDP, well beyond the intrinsic bounds one has in [4]: remarkably enough, the outcome still provides quantizations of a coisotropic subgroup, or a Poisson quotient, closely related to the initial subobject in terms of Poisson duality again. From this point of view our functors will provide sort of a Galois' reciprocity, in which coisotropic objects play the rôle of closed ones.

In the end, we present a few explicit examples to describe all these features; in particular, this shows how explicit computations may be carried out to produce new examples of quantum homogeneous spaces (actually, of Poisson quotients).

Some of the proofs involved in this global QDP are nothing but simple analogues of those given in [4]: in such cases we will simply indicate to the reader the relevant statements in our previous work. Our main purpose instead will be to stress where some different behaviour shows up.

2 General Theory

To collect in the present section some classical material about Poisson geometry for groups and homogeneous spaces. Everything is standard, we just need to fix the main notions and notations we shall deal with.

2.1 Subgroups and homogeneous spaces

Let G be a complex affine algebraic group and let $\mathfrak{g}$ be its tangent Lie algebra. Let us denote by $F[G]$ its algebra of regular functions and by $U(\mathfrak{g})$ its universal enveloping algebra. Both such algebras are Hopf algebras, and there exists a natural pairing of Hopf algebras between them, given by evaluation of differential operators onto functions. This pairing is perfect if and only if G is connected, which we will always assume in what follows.

A *real form* of either G or $\mathfrak{g}$ is given once Hopf $*$–algebras structure, dual to each other, are considered on both $F[G]$ and $U(\mathfrak{g})$. Thus by *real algebraic group* we will always mean a complex algebraic group with a suitable $*$–structure.

A subgroup K of G will always be considered as Zariski–closed and algebraic. For any such subgroup, the quotient G/K is an algebraic left homogeneous G–space, which is quasi-projective as an algebraic variety. Given an algebraic left homogeneous G–space M and choosing $m \in M$, the stabilizer subgroup K_m will be a closed algebraic subgroup of G such that: $G\big/K_m \simeq M$. Changing point will change the stabilizer within a conjugacy class.

We would like to describe the subgroup K, or the homogeneous space G/K, through either an algebraic subset of $F[G]$ — to which we will refer as a *global* coding — or an algebraic subset of $U(\mathfrak{g})$ — to which we will refer as a *local* coding. The complete picture is the following:

subgroup K

- *(local)* letting $\mathfrak{k} = \mathrm{Lie}(K)$ we can consider its enveloping algebra $U(\mathfrak{k})$ which is a Hopf subalgebra of $U(\mathfrak{g})$. We will let $\mathfrak{C} \equiv \mathfrak{C}(K) := U(\mathfrak{k})$;
- *(global)* functions which are 0 on K form a Hopf ideal $\mathcal{I} \equiv \mathcal{I}(K)$ inside $F[G]$, such that $F[K] \simeq F[G]\big/\mathcal{I}$.

homogeneous space G/K

- *(local)* let $\mathfrak{I} \equiv \mathfrak{I}(K) = U(\mathfrak{g}) \cdot \mathfrak{k}$. This is a left ideal and two-sided coideal in $U(\mathfrak{g})$, and $U(\mathfrak{g})/\mathfrak{I}$ is the set of left–invariant differential operators on G/K.
- *(global)* functions on the homogeneous space G/K may be identified with K–invariant functions on G. We will let $\mathcal{C} = \mathcal{C}(K) = F[G]^K$; this is a subalgebra and left coideal in $F[G]$.

Let us introduce, for any Hopf algebra $\mathcal{H}$ the following notations: $\leq^1$ will stand for unital subalgebra, $\trianglelefteq$ for two-sided ideal, $\trianglelefteq_l$ for left ideal and similarly $\dot{\leq}$ will stand for subcoalgebra, $\dot{\trianglelefteq}$ for two-sided coideal and $\dot{\trianglelefteq}_l$ for left coideal. When the same symbols will be decorated by a subindex referring to a specific algebraic structure their meaning should be modified accordingly, for example $\trianglelefteq_{\mathcal{H}}$ will stand for Hopf ideal and $\leq_{\mathcal{H}}$ for Hopf subalgebra. With such notations, to any subgroup K of G there is associated one of the following algebraic objects:

$$(a)\ \mathcal{I} \trianglelefteq_{\mathcal{H}} F[G] \quad (b)\ \mathcal{C} \leq^1 \dot{\trianglelefteq}_l\, F[G] \quad (c)\ \mathfrak{I} \trianglelefteq_l\ \dot{\trianglelefteq}\, U(\mathfrak{g}) \quad (d)\ \mathfrak{C} \leq_{\mathcal{H}} U(\mathfrak{g}) \qquad (2.1)$$

In the real case, additional requirements involving the $*$ structure and the antipode S have to be considered together with (2.1):

$$(a)\ \mathcal{I}^* = \mathcal{I} \qquad (b)\ S(\mathcal{C})^* = \mathcal{C} \qquad (c)\ S(\mathfrak{I})^* = \mathfrak{I} \qquad (d)\ \mathfrak{C}^* = \mathfrak{C} \qquad (2.2)$$

In the connected case algebraic objects of type $\mathcal{I}$, $\mathfrak{I}$ and $\mathfrak{C}$ in (2.1) are enough to reconstruct either K or $G\big/K$:

$$K \;=\; \mathrm{Spec}\big(F[G]/\mathcal{I}\big) \;=\; \exp\big(\mathrm{Prim}(\mathfrak{C})\big) \;=\; \exp\big(\mathrm{Prim}(\mathfrak{I})\big)$$

where $\mathrm{Prim}(X)$ denotes the set of primitive elements of a bialgebra X). In contrast, $\mathcal{C}$ might be not enough to reconstruct K, due to lack of enough global algebraic functions; this happens, for example,when $G\big/K$ is projective and therefore $\mathcal{C}(K) = \mathbb{C}$. Any group K which can be reconstructed from its associated $\mathcal{C}$is called *observable*: we shall now make this notion more precise.

Let us call τ the map that to any subgroup K associates the algebra of invariant functions $F[G]^K$ and let us call σ the map that to any subalgebra of $F[G]$ associates its stabilizer $\sigma(A) = \big\{ g \in G \,\big|\, g \cdot f = f \;\; \forall f \in F[G] \big\}$. These two maps are obviously inclusion–reversing. Furthermore they establish what is also known as a *simple Galois correspondence*, i.e. for any subgroup K and subalgebra A

$$(\sigma \circ \tau)(K) \supseteq K \,, \qquad (\tau \circ \sigma)(A) \subseteq A$$

so that $(\tau \circ \sigma \circ \tau)\,(K) = K$, $(\sigma \circ \tau \circ \sigma)\,(A) = A$. A subgroup K of G such that $(\sigma \circ \tau)\,(K) = K$ is said to be *observable*: this means exactly that such a subgroup can be fully recovered from its algebra of invariant functions $\tau(K)$. If K is any subgroup, then $\widehat{K} := (\sigma \circ \tau)\,(K)$ is the smallest observable subgroup containg K; we will call it the *observable hull* of K. Remark then that $\mathcal{C}(K) = \mathcal{C}(\widehat{K})$.

The following fact (together with many properties of observable subgroups), which gives a characterization of observable subgroups in purely geometrical terms, may be found in [11]:

Fact: a subgroup K of G is observable if and only if G/K is quasi–affine.

Let us now clarify how to pass from algebraic objects directly associated to subgroups to those corresponding to homogeneous spaces. Let H be a Hopf algebra, with counit ε. For any submodule $M \subseteq H$ define

$$M^+ := M \cap Ker(\varepsilon) \;, \quad H^{coM} := \big\{ y \in H \,\big|\, (\delta(y) - y \otimes 1) \in H \otimes M \big\} \qquad (2.3)$$

Let C be a (unital) subalgebra and left coideal of H and define $\Psi(C) = H \cdot C^+$. Then $\Psi(C)$ is a left ideal and two-sided coideal in H. Conversely, let I be a left ideal and two-sided coideal in H and define $\Phi(I) := H^{coI}$. Then $\Phi(I)$ is a unital subalgebra and left coideal in H. Also, this pair of maps (Φ, Ψ) defines a simple Galois correspondence, that is to say *(a)* Ψ is inclusion-preserving, Φ is inclusion-reversing;

(b) $(\Phi \circ \Psi)\,(C) \supseteq C\ , \quad (\Psi \circ \Phi)\,(I) \subseteq I\ ;$

(c) $\Phi \circ \Psi \circ \Phi = \Phi\ , \quad \Psi \circ \Phi \circ \Psi = \Psi\ .$

(where the third property follows from the previous ones; see [17, 18, 19] for further details).

Let now K be a subgroup of G and let $\mathcal{I}$, $\mathcal{C}$, $\mathfrak{I}$, $\mathfrak{C}$ the corresponding algebraic objects as described in (2.1). We can thus establish the following relations among them:

subgroup vs. homogeneous space: objects directly related to the subgroup (i.e. $\mathcal{I}$ and $\mathfrak{C}$) and objects directly related to the homogeneous space (i.e. $\mathcal{C}$ and $\mathfrak{I}$) are linked by Ψ and Φ as follows:

$$\mathfrak{I} = \Psi(\mathfrak{C})\,, \quad \mathfrak{C} = \Phi(\mathfrak{I})\,, \quad \mathcal{I} \supseteq \Psi(\mathcal{C})\,, \quad \mathcal{C} = \Psi(\mathcal{I}) \tag{2.4}$$

In particular, K is observable if and only if $\mathcal{I} = \Psi(\mathcal{C})$; on the other hand, we have in general $\Psi(\mathcal{C}(K)) = \mathcal{I}(\widehat{K})$.

orthogonality with respect to the natural pairing between $F[G]$ and $U(\mathfrak{g})$: this is expressed by

$$\mathcal{I} = \mathfrak{C}^{\perp}\,, \quad \mathfrak{C} = \mathcal{I}^{\perp}\,, \quad \mathcal{C} = \mathfrak{I}^{\perp}\,, \quad \mathfrak{I} \subseteq \mathcal{C}^{\perp}\,. \tag{2.5}$$

In particular, K is observable if and only if $gerI = \mathcal{C}^{\perp}$; on the other hand, we have in general $\mathcal{C}(K)^{\perp} = \mathfrak{I}(\widehat{K})$.

Let us also remark that orthogonality intertwines the local and global description.

2.2 Poisson subgroups and Poisson quotients

Let us now assume that G is endowed with a complex Poisson group structure corresponding to a Lie bialgebra structure on $\mathfrak{g}$, whose Lie cobracket is denoted $\delta : \mathfrak{g} \longrightarrow \mathfrak{g} \wedge \mathfrak{g}$. At the Hopf algebra level this means that $F[G]$ is a Poisson–Hopf algebra and $U(\mathfrak{g})$ a co-Poisson Hopf algebra, in such a way that the duality pairing is compatible with these additional structures (see [3] for basic definitions). Let us recall that the linear dual $\mathfrak{g}^*$ inherits a Lie algebra structure; on the other hand, it has a natural Lie coalgebra structure, whose cobracket $\delta : \mathfrak{g}^* \longrightarrow \mathfrak{g}^* \wedge \mathfrak{g}^*$ is the dual map to the Lie bracket of $\mathfrak{g}$. Altogether, this makes $\mathfrak{g}^*$ into a Lie bialgebra, which said to be *dual* to $\mathfrak{g}$. Therefore, there exist Poisson groups whose tangent Lie bialgebra is $\mathfrak{g}^*$; we will assume one such connected group is fixed, we will denote it with G^* and call it the dual Poisson group of G. In the real case the involution in $F[G]$ is a Poisson algebra antimorphism and the one in $U(\mathfrak{g})$ is a co-Poisson algebra antimorphism.

A closed subgroup K of G is called *coisotropic* if its defining ideal $\mathcal{I}(K)$ is a Poisson subalgebra, while it is called a *Poisson subgroup* if $\mathcal{I}(K)$ is a Poisson ideal, the latter condition being equivalent to $K \hookrightarrow G$ being a Poisson map. Connected coisotropic subgroups can be characterized, at an infinitesimal level, by one of the following conditions on $\mathfrak{k} \subseteq \mathfrak{g}$:

(C-i) $\delta(\mathfrak{k}) \subseteq \mathfrak{k} \wedge \mathfrak{g}$, that is $\mathfrak{k}$ is a Lie coideal in $\mathfrak{g}$,
(C-ii) $\mathfrak{k}^\perp$ is a Lie subalgebra of $\mathfrak{g}^*$,

while analogous characterizations of Poisson subgroups correspond to $\mathfrak{k}$ being a Lie subcoalgebra or $\mathfrak{k}^\perp$ being a Lie ideal.

The most important features of coisotropic subgroups, in this setting, is the fact that G/K naturally inherits a Poisson structure from that of G. A Poisson manifold $(M\,\omega_M)$ is a *Poisson homogeneous G–space* if there exists a smooth, homogeneous G–action $\phi : G \times M \to M$ which is a Poisson map (w.r. to the product Poisson structure on the domain). We will say that (M, ω_M) is a *Poisson quotient* if it verifies one of the following equivalent conditions ([21]):

(P-i) there exists $x_0 \in M$ whose stabilizer G_{x_0} is coisotropic in G ;
(P-ii) there exists $x_0 \in M$ such that $\phi_{x_0} : G \to M$, $\phi(x_0, g) = \phi(x, g)$, is a Poisson map ;
(P-iii) there exists $x_0 \in M$ such that $\omega_M(x_0) = 0$.

It is important to remark here that inside the same conjugacy class of subgroups of G there may be subgroups which are Poisson, coisotropic, non coisotropic. Therefore, on the same homogeneous space, there maybe many Poisson homogeneous structures, some of which are Poisson quotients and some of which are not.

For a fixed connected subgroup K of a Poisson group G, with Lie algebra $\mathfrak{k}$, one can consider the following descriptions in terms of the Poisson Hopf algebra $F[G]$ or of the co-Poisson Hopf algebra $U(\mathfrak{g})$:

$$\mathcal{I} \leq_{\mathcal{P}} F[G] \, , \qquad \mathcal{C} \leq_{\mathcal{P}} F[G] \tag{2.6}$$

$$\mathfrak{I} \mathrel{\dot{\trianglelefteq}}_{\mathcal{P}} U(\mathfrak{g}) \, , \qquad \mathfrak{C} \mathrel{\dot{\trianglelefteq}}_{\mathcal{P}} U(\mathfrak{g}) \tag{2.7}$$

where on first line we have global conditions and on second line local ones. Conversely each one of these conditions imply coisotropy of G with the exception of the condition on $\mathcal{C}$, which implies only that the observable hull $\widehat{K}$ is coisotropic. Therefore a connected, observable, coisotropic subgroup of G is identified by one of the following algebraic objects:

$$\mathcal{I} \trianglelefteq_{\mathcal{H}} \leq_{\mathcal{P}} F[G] \, , \qquad \mathcal{C} \leq^1 \dot{\trianglelefteq}_l \leq_{\mathcal{P}} F[G] \tag{2.8}$$

$$\mathfrak{I} \trianglelefteq_l \dot{\trianglelefteq} \, \dot{\trianglelefteq}_{\mathcal{P}} U(\mathfrak{g}) \, , \qquad \mathfrak{C} \leq_{\mathcal{H}} \dot{\trianglelefteq}_{\mathcal{P}} U(\mathfrak{g}) \tag{2.9}$$

Thanks to selfduality in the notion of Lie bialgebra, with any Poisson group there is associated a natural *Poisson dual*, which is fundamental in the QDP. If QDP has to be extended to coisotropic subgroups a suitable duality at the Poisson level is needed.

Definition 2.1. *Let G be a Poisson group and G^* a fixed Poisson dual.*

1. *If K is coisotropic in G we call* complementary dual of K *the unique connected subgroup $K^\perp$ in G^* such that* $\mathrm{Lie}(K^\perp) = \mathfrak{k}^\perp$.
2. *If M is a Poisson quotient and $M \simeq G\big/K_M$ we call* complementary dual of M *the Poisson G^*–quotient $M^\perp := G^*\big/K_M^\perp$.*
3. *For any subgroup H of G we call* coisotropic interior of H *the unique maximal, closed, connected, coisotropic subgroup $\overset{\circ}{H}$ of G contained in H .*

Remarks:

1. The complementary dual of a coisotropic subgroup is, trivially, a coisotropic subgroup whose complementary dual is the one we started with. Similarly the complementary dual of a Poisson quotient is a Poisson quotient (by definition) and taking twice the complementary dual brings back to the original Poisson quotient.
2. The coisotropic interior may be characterized, at an algebraic level, as the unique closed subgroup whose Lie algebra is maximal between Lie subalgebras of $\mathfrak{h}$ which are Lie coideals in $\mathfrak{g}$.

Proposition 2.2. *Let K be any subgroup of G and let $K^{\langle\perp\rangle} := \big\langle \exp(\mathfrak{k}^\perp) \big\rangle$ be the closed, connected, subgroup of G^* generated by $\exp(\mathfrak{k}^\perp)$. Then:*

(a) the Lie algebra $\mathfrak{k}^{\langle\perp\rangle}$ of $K^{\langle\perp\rangle}$ is the Lie subalgebra of $\mathfrak{g}^*$ generated by $\mathfrak{k}^\perp$;

(b) $\mathfrak{k}^{\langle\perp\rangle}$ is a Lie coideal of $\mathfrak{g}^*$, hence $K^{\langle\perp\rangle}$ is a coisotropic subgroup of G^*;

(c) $K^{\langle\perp\rangle} = (\overset{\circ}{K})^\perp$; in particular if K is coisotropic then $K^{\langle\perp\rangle} = K^\perp$;

(d) $(K^{\langle\perp\rangle})^{\langle\perp\rangle} = \overset{\circ}{K}$ and K is coisotropic if and only if $(K^{\langle\perp\rangle})^{\langle\perp\rangle} = K$.

Proof. Part *(a)* is trivial. As for *(b)*, since $\mathfrak{k} = (\mathfrak{k}^\perp)^\perp$ is a Lie subalgebra of $\mathfrak{g}$, we have that $\mathfrak{k}^\perp$ is a Lie coideal in $\mathfrak{g}^*$: therefore, due to the identity

$$\delta\big([x,y]\big) \;=\; \sum_{[y]} \big([x,y_{[1]}] \otimes y_{[2]} + y_{[1]} \otimes [x,y_{[2]}]\big) + \sum_{[x]} \big([x_{[1]},y] \otimes x_{[2]} + x_{[1]} \otimes [x_{[2]},y]\big)$$

(where $\delta(z) = \sum_{[z]} z_{[1]} \otimes z_{[2]}$ for $z \in \mathfrak{g}^*$), the Lie subalgebra $\langle \mathfrak{k}^\perp \rangle$ of $\mathfrak{g}^*$ generated by $\mathfrak{k}^\perp$ is a Lie coideal too. It follows then by claim *(a)* that $K^{\langle\perp\rangle}$ is coisotropic. Thus *(b)* is proved.

As for part *(c)* we have

$$\big(\mathfrak{k}^{\langle\perp\rangle}\big)^\perp \;=\; \langle \mathfrak{k}^\perp \rangle^\perp \;=\; \Bigg(\bigcap_{\substack{\mathfrak{h} \leq_{\mathcal{L}} \mathfrak{g}^* \\ \mathfrak{h} \supseteq \mathfrak{k}^\perp}} \mathfrak{h} \Bigg)^{\perp} \;=\; \sum_{\substack{\mathfrak{h} \leq_{\mathcal{L}} \mathfrak{g}^* \\ \mathfrak{h} \supseteq \mathfrak{k}^\perp}} \mathfrak{h} \;=\; \sum_{\substack{\mathfrak{f} \,\dot{\trianglelefteq}_{\mathcal{L}}\, \mathfrak{g} \\ \mathfrak{f} \supseteq \mathfrak{k}}} \mathfrak{f} \;= \overset{\circ}{\mathfrak{k}}$$

(with $\leq_{\mathcal{L}}$ meaning “Lie subalgebra” and $\dot{\trianglelefteq}_{\mathcal{L}}$ meaning “Lie coideal”) where $\overset{\circ}{\mathfrak{k}}$ is exactly the maximal Lie subalgebra and Lie coideal of $\mathfrak{g}$ contained in $\mathfrak{k}$. To be precise, this last statement follows from the above formula for $\delta\big([x,y]\big)$, since that formula implies that the Lie subalgebra generated by a family of Lie coideals is still a Lie coideal.

Now $\mathring{\mathfrak{k}} = \mathrm{Lie}(\mathring{K})$, so $\mathrm{Lie}\big(K^{\langle\perp\rangle}\big) = \mathfrak{k}^{\langle\perp\rangle} = \Big(\big(\mathfrak{k}^{\langle\perp\rangle} \big)^{\perp} \Big)^{\perp} = \big(\mathring{\mathfrak{k}} \big)^{\perp} = \mathrm{Lie}(\mathring{K})^{\perp}$ implies $K^{\langle\perp\rangle} = (\mathring{\mathfrak{k}})^{\perp}$ as we wished to prove. If, in addition, K is coisotropic then, obviously, $K^{\langle\perp\rangle} = K$. All other statements follow easily. $\square$

3 Strict, proper, weak quantizations

The purpose of this section is to fix some terminology concerning the meaning of the word "quantization" and to describe some possible ways of quantizing a homogeneous space. We set the algebraic machinery needed for talking of "quantization" and "specialization": these notions must be carefully specified before approaching the construction of Drinfeld's functors. Roughly speaking, a quantum group is a Hopf algebra H_q depending on a parameter, say q, such that at some special value $q = q_0$ one gets either $H_{q_0} = F[G]$ or $H_{q_0} = U(\mathfrak{g})$, for some Poisson group G or Lie bialgebra $\mathfrak{g}$.

Let q be an indeterminate, $\mathbb{C}\big[q, q^{-1}\big]$ the ring of complex-valued Laurent polynomials in q, and $\mathbb{C}(q)$ the field of complex-valued rational functions in q. Denote by $\mathcal{HA}$ the category of all Hopf algebras over $\mathbb{C}\big[q, q^{-1}\big]$ which are torsion-free as $\mathbb{C}\big[q, q^{-1}\big]$–modules.

Given a Hopf $\mathbb{C}(q)$–algebra H in $\mathcal{HA}_F$, a subset $\overline{H} \subseteq H$ is called *a* $\mathbb{C}\big[q, q^{-1}\big]$*–integral form* (or simply *a* $\mathbb{C}\big[q, q^{-1}\big]$*–form*) if it is a $\mathbb{C}\big[q, q^{-1}\big]$–Hopf subalgebra of H and $H_F := \mathbb{C}(q) \otimes_{\mathbb{C}[q,q^{-1}]} \overline{H} = H$. Note then that $\overline{H}$ is torsion-free as a $\mathbb{C}\big[q, q^{-1}\big]$–module, hence $\overline{H} \in \mathcal{HA}$.

For any $\mathbb{C}\big[q, q^{-1}\big]$–module M, we set $M_1 := M \big/ (q-1) M = \mathbb{C} \otimes_{\mathbb{C}[q,q^{-1}]} M$: this is a $\mathbb{C}$–module (via $\mathbb{C}\big[q, q^{-1}\big] \rightarrow \mathbb{C}\big[q, q^{-1}\big] \Big/ (q-1) = \mathbb{C}$), called *specialization of* M *at* $q = 1$.

Given two $\mathbb{C}(q)$–modules A and B and a $\mathbb{C}(q)$–bilinear pairing $A \times B \longrightarrow F$, for any $\mathbb{C}\big[q, q^{-1}\big]$–submodule $A_\times \subseteq A$ or $B_\times \subseteq B$ we set:

$$A_\times^{\bullet} \;:=\; \Big\{ b \in B \,\Big|\, \big\langle A_\times, b \big\rangle \subseteq \mathbb{C}\big[q, q^{-1}\big] \Big\}\,, \tag{3.1}$$

$$A_\times^{\perp_q} \;:=\; \Big\{ b \in B \,\Big|\, \big\langle A_\times, b \big\rangle \subseteq (q-1)\mathbb{C}\big[q, q^{-1}\big] \Big\} \tag{3.2}$$

In such a setting, we call $A_\times^{\bullet}$ *the* $\mathbb{C}\big[q, q^{-1}\big]$*–dual of* $A_\times$.

We will call *quantized universal enveloping algebra* (or, in short, QUEA) any $U_q \in \mathcal{HA}$ such that $U_1 := (U_q)_1$ is isomorphic to $U(\mathfrak{g})$ for some Lie algebra $\mathfrak{g}$, and we will call *quantized function algebra* any $F_q \in \mathcal{HA}$ such that $F_1 := \big(F_q\big)_1$ is isomorphic to $F[G]$ for some connected algebraic group G and, in addition,

$$\bigcap_{n \geq 0} (q-1)^n F_q \;=\; \bigcap_{n \geq 0} \big((q-1) F_q \,+\, Ker(\epsilon_{F_q}) \big)^n$$

We will add the specification that such quantum algebras are *real* whenever the starting object is a $*$–Hopf algebra. As a matter of notation, we write

$$\mathbb{U}_q \;:=\; \mathbb{C}(q) \otimes_{\mathbb{C}[q,q^{-1}]} U_q \quad , \qquad \mathbb{F}_q \;:=\; \mathbb{C}(q) \otimes_{\mathbb{C}[q,q^{-1}]} F_q \quad .$$

Whenever U_q is a (real) QUEA, its specialization U_1 is a (real) co–Poisson Hopf algebra so that $\mathfrak{g}$ is in fact a (real) Lie bialgebra. Similarly, for any (real) QFA F_q the specialization F_1 is a (real) Poisson-Hopf algebra and therefore G is a (real) Poisson group (see [3] for details).

On occasions it is useful to consider simultaneous quantizations of both the universal enveloping algebra and the function algebra, or, in a larger generality, of a pair of dual Hopf algebra. Let $H, K \in \mathcal{HA}$ and assume that there exists a pairing of Hopf algebras $H \times K \to \mathbb{C}[q, q^{-1}]$. The same symbol will denote its scalar extension to a pairing between with values in $\mathbb{C}(q)$. If the pairing is such that

(a) $H = K^\bullet$, $K = H^\bullet$ (notation of (3.1)) w.r.t. the pairing $\mathbb{H} \times \mathbb{K} \to \mathbb{C}(q)$, for $\mathbb{H} := \mathbb{C}(q) \otimes_{\mathbb{C}[q,q^{-1}]} H$, $\mathbb{K} := \mathbb{C}(q) \otimes_{\mathbb{C}[q,q^{-1}]} K$, induced from $H \times K \to \mathbb{C}(q)$;

(b) the Hopf pairing $H_1 \times K_1 \to \mathbb{C}$ given by specialization at $q = 1$ is perfect.

Then these requirements will imply that the initial pairing is perfect too and we will say that H and K are dual to each other. In the case in which $H = U_q(\mathfrak{g})$ is a QUEA and $K = F_q[G]$ is a QFA, if the specialized pairing at 1 is the natural pairing between $U(\mathfrak{g})$ and $F[G]$ we will say that *the pair* $(U_q(\mathfrak{g}), F_q[G])$ *is a double quantization of* $(G, \mathfrak{g})$.

Let us now move to the case in which G is a Poisson group and K a subgroup. We want to define a reasonable quantization of K and of the corresponding homogeneous space $G \big/ K$. There is a standard way to implement this, with the additional requirement that K is a coisotropic subgroup of G .

Definition 3.1. *Let* $F_q[G]$ *and* $U_q(\mathfrak{g})$ *be a QFA and a QUEA for* G *and* $\mathfrak{g}$ *and let*

$$\pi_{F_q} \colon\, F_q[G] \twoheadrightarrow F_q[G] \big/ (q-1) F_q[G] \,\cong\, F[G]$$

$$\pi_{U_q} \colon U_q(\mathfrak{g}) \twoheadrightarrow U_q(\mathfrak{g}) \big/ (q-1)\, U_q(\mathfrak{g}) \cong U(\mathfrak{g})$$

be the specialization maps. Let $\mathcal{I}$, $\mathcal{C}$, $\mathfrak{I}$ *and* $\mathfrak{C}$ *be the algebraic objects associated with the subgroup* K *of* G *(see 2.1). We call "strict quantization" (and sometimes we shall drop the adjective "strict") of each of them any object* $\mathcal{I}_q$, $\mathcal{C}_q$, $\mathfrak{I}_q$ *or* $\mathfrak{C}_q$ *respectively, such that*

$$\begin{array}{llll} (a) & \mathcal{I}_q \trianglelefteq_\ell \dot{\trianglelefteq}\, F_q[G] \ , & \pi_{F_q}(\mathcal{I}_q) \,=\, \mathcal{I} \ , & \pi_{F_q}(\mathcal{I}_q) \,\cong\, \mathcal{I}_q \big/ (q-1)\, \mathcal{I}_q \\ (b) & \mathcal{C}_q \leq^1 \dot{\trianglelefteq}_\ell\, F_q[G] \ , & \pi_{F_q}(\mathcal{C}_q) \,=\, \mathcal{C} \ , & \pi_{F_q}(\mathcal{C}_q) \,\cong\, \mathcal{C}_q \big/ (q-1)\, \mathcal{C}_q \\ (c) & \mathfrak{I}_q \trianglelefteq_\ell \dot{\trianglelefteq}\, U_q(\mathfrak{g}) \ , & \pi_{U_q}(\mathfrak{I}_q) \,=\, \mathfrak{I} \ , & \pi_{U_q}(\mathfrak{I}_q) \,\cong\, \mathfrak{I}_q \big/ (q-1)\, \mathfrak{I}_q \\ (d) & \mathfrak{C}_q \leq^1 \dot{\trianglelefteq}_\ell\, U_q(\mathfrak{g}) \ , & \pi_{U_q}(\mathfrak{C}_q) \,=\, \mathfrak{C} \ , & \pi_{U_q}(\mathfrak{C}_q) \,\cong\, \mathfrak{C}_q \big/ (q-1)\, \mathfrak{C}_q \end{array} \tag{3.3}$$

To explain this definition let us start by considering the first two conditions in each line of (3.3).

a) A left ideal and two-sided coideal in a QFA quantizes the Hopf ideal of functions which are zero on a *subgroup*;

b) a left coideal subalgebra in a QFA quantizes the algebra of invariant functions on a *homogeneous space*;

c) a left ideal and two-sided coideal in a QUEA quantizes the infinitesimal algebra on a *homogeneous space*;

d) a left coideal subalgebra in a QUEA quantizes the universal enveloping subalgebra of a *subgroup*.

It has to be noted here that $\mathcal{C}_q$, for reasons explained in Proposition 2.4, has to be seen as a quantization of the observable hull $\widehat{K}$ rather than of K itself: this follows from a classical feature, because from $\mathcal{C}$ we can only reconstruct $\widehat{K}$, not K.

Let us now be more precise about the last condition in the previous definition. By asking $\mathcal{I}_q \big/ (q-1)\mathcal{I}_q \cong \pi_{F_q}(\mathcal{I}_q) = \mathcal{I}$ we mean the following: the specialization map sends $\mathcal{I}_q$ inside $F[G]$. This map factors through $\mathcal{I}_q \big/ (q-1)\mathcal{I}_q$; in addition, we require that the induced map $\mathcal{I}_q \big/ (q-1)\mathcal{I}_q \longrightarrow F[G]$ be a bijection on $\mathcal{I}$. Of course this bijection will respect the whole Hopf structure, since π_{F_q} does. Now, since

$$\pi_{F_q}(\mathcal{I}_q) \, = \, \mathcal{I}_q \Big/ \big(\mathcal{I}_q \cap (q-1) F_q[G] \big)$$

we also have that this property may be equivalently rephrased by saying that $\mathcal{I}_q \cap (q-1) F_q[G] = (q-1)\mathcal{I}_q$. The previous discussions may be repeted unaltered for all four algebraic objects under consideration. An equivalent definition of strict quantizations is therefore the following:

$$\begin{array}{lllll} (a) & \mathcal{I}_q \trianglelefteq_\ell \dot{\trianglelefteq} \, F_q[G] \ , & \pi_{F_q}(\mathcal{I}_q) \, = \, \mathcal{I} \ , & \mathcal{I}_q \cap (q-1) F_q[G] \, = \, (q-1)\, \mathcal{I}_q \\ (b) & \mathcal{C}_q \leq^1 \dot{\trianglelefteq}_\ell \, F_q[G] \ , & \pi_{F_q}(\mathcal{C}_q) \, = \, \mathcal{C} \ , & \mathcal{C}_q \cap (q-1) F_q[G] \, = \, (q-1)\, \mathcal{C}_q \\ (c) & \mathfrak{I}_q \trianglelefteq_\ell \dot{\trianglelefteq} \, U_q(\mathfrak{g}) \ , & \pi_{U_q}(\mathfrak{I}_q) \, = \, \mathfrak{I} \ , & \mathfrak{I}_q \cap (q-1) U_q(\mathfrak{g}) \, = \, (q-1)\, \mathfrak{I}_q \\ (d) & \mathfrak{C}_q \leq^1 \dot{\trianglelefteq}_\ell \, U_q(\mathfrak{g}) \ , & \pi_{U_q}(\mathfrak{C}_q) \, = \, \mathfrak{C} \ , & \mathfrak{C}_q \cap (q-1) U_q(\mathfrak{g}) \, = \, (q-1)\, \mathfrak{C}_q \end{array} \tag{3.4}$$

The purpose of the last condition — which is often mentioned by saying that $\mathfrak{C}_q$ is a *flat* quantization (typically, in the literature on deformation quantization) — should be clear: indeed, removing it means losing any control on what is contained, in quantization, inside the kernel of the specialization map.

Although the just mentioned notion of quantization appears to be, in many respect, the "correct" one, another notion of quantization naturally appears when one has to deal with quantum duality principle.

Definition 3.2. *Let $F_q[G]$ and $U_q(\mathfrak{g})$ be a QFA and a QUEA for G and $\mathfrak{g}$ and let*

$$\pi_{F_q} \colon \, F_q[G] \relbar\joinrel\twoheadrightarrow F_q[G] \Big/ (q-1) F_q[G] \, \cong \, F[G]$$

$$\pi_{U_q} \colon U_q(\mathfrak{g}) \relbar\joinrel\twoheadrightarrow U_q(\mathfrak{g}) \Big/ (q-1)\, U_q(\mathfrak{g}) \, \cong \, U(\mathfrak{g})$$

be the specialization maps. Let $\nabla := \Delta - \Delta^{op}$. Let $\mathcal{I}$, $\mathcal{C}$, $\mathfrak{I}$ and $\mathfrak{C}$ be the algebraic objects associated with the subgroup K of G (see 2.1). We call "proper quantization" of each of them any object $\mathcal{I}_q$, $\mathcal{C}_q$, $\mathfrak{I}_q$ or $\mathfrak{C}_q$ respectively, such that

$$\begin{array}{lllll} (a) & \mathcal{I}_q \trianglelefteq_\ell \dot{\trianglelefteq} \, F_q[G] \ , & \pi_{F_q}(\mathcal{I}_q) \, = \, \mathcal{I} \ , & \big[\mathcal{I}_q \, , \mathcal{I}_q \big] \, \subseteq \, (q-1)\, \mathcal{I}_q \\ (b) & \mathcal{C}_q \leq^1 \dot{\trianglelefteq}_\ell \, F_q[G] \ , & \pi_{F_q}(\mathcal{C}_q) \, = \, \mathcal{C} \ , & \big[\mathcal{C}_q \, , \mathcal{C}_q \big] \, \subseteq \, (q-1)\, \mathcal{C}_q \\ (c) & \mathfrak{I}_q \trianglelefteq_\ell \dot{\trianglelefteq} \, U_q(\mathfrak{g}) \ , & \pi_{U_q}(\mathfrak{I}_q) \, = \, \mathfrak{I} \ , & \nabla(\,\mathfrak{I}_q) \, \subseteq \, (q-1)\, U_q(\mathfrak{g}) \wedge \mathfrak{I}_q \\ (d) & \mathfrak{C}_q \leq^1 \dot{\trianglelefteq}_\ell \, U_q(\mathfrak{g}) \ , & \pi_{U_q}(\mathfrak{C}_q) \, = \, \mathfrak{C} \ , & \nabla(\mathfrak{C}_q) \, \subseteq \, (q-1)\, U_q(\mathfrak{g}) \wedge \mathfrak{C}_q \end{array} \tag{3.5}$$

The link between these two notions of quantization is the following:

Lemma 3.3. *Any strict quantization is a proper quantization.*

Proof. This is an easy consequence of definitions. Indeed, let K be a subgroup of G. If $\mathcal{I}_q := \mathcal{I}\big(\widehat{K}\big)$ is any strict quantization of $\mathcal{I}(K)$, we have

$$\mathcal{I}_q \cap (q-1)\, F_q \;=\; (q-1)\, \mathcal{I}_q$$

by assumption, and moreover $\big[F_q\,, F_q\big] \subseteq (q-1)\, F_q$. Then

$$\big[\mathcal{I}_q\,, \mathcal{I}_q\big] \;\subseteq\; \mathcal{I}_q \cap \big[F_q\,, F_q\big] \;\subseteq\; \mathcal{I}_q \cap (q-1)\, F_q \;=\; (q-1)\, \mathcal{I}_q$$

thus $\big[\mathcal{I}_q\,, \mathcal{I}_q\big] \subseteq (q-1)\, \mathcal{I}_q$, i.e. $\mathcal{I}_q$ is proper. A similar argument works for quantizations of type $\mathcal{C}_q(K)$. Also, if $\mathfrak{I}_q(K)$ is any strict quantization of $\mathfrak{I}(K)$, then we have $\mathfrak{I}_q \cap (q-1)\, U_q \;=\; (q-1)\, \mathfrak{I}_q$ by assumption, and moreover $\nabla(U_q) \;\subseteq\; (q-1)\, U_q^{\wedge 2}$. Then

$$\nabla(\,\mathfrak{I}_q) \;\subseteq\; \big(U_q \wedge \mathfrak{I}_q\big) \cap \nabla(U_q) \;\subseteq\; \big(U_q \wedge \mathfrak{I}_q\big) \cap (q-1)\, U_q^{\wedge 2} \;\subseteq\; (q-1)\, U_q \wedge \mathfrak{I}_q$$

so that $\mathfrak{I}_q$ is proper. A similar argument works for quantizations of type $\mathfrak{C}_q(K)$. $\square$

Remark 3.4. *The converse to Lemma 3.3 here above is false.*

Indeed, there exist quantizations (of subgroups / homogeneous spaces) which are proper but *not* strict: we present an explicit example — of type $\mathcal{C}_q$ — in Subsection 6.3 later on.

The following statement clarifies why such definitions are reasonable in the restricted case of coisotropic subgroups (this result can be traced back to [16], where it is mentioned as *coisotropic creed*).

Proposition 3.5. *Let K be a subgroup of G and assume a proper quantization of it exists. Then K is coisotropic or, in case the quantization is $\mathcal{C}_q$, its observable hull $\widehat{K}$ is coisotropic.*

Proof. Assume $\mathcal{I}_q$ exists. Let $f, g \in \mathcal{I}$, and let $\varphi, \gamma \in \mathcal{I}_q$ with $\pi_{F_q}(\varphi) = f$, $\pi_{F_q}(\gamma) = g$. Then by definition $\{f, g\} = \pi_{F_q}\big((q-1)^{-1}[\varphi, \gamma]\big)$. But

$$[\varphi, \gamma] \in \big[\mathcal{I}_q\,, \mathcal{I}_q\big] \subseteq (q-1)\, \mathcal{I}_q$$

by assumption, hence $(q-1)^{-1}[\varphi, \gamma] \in \mathcal{I}_q$, thus $\{f, g\} = \pi_{F_q}\big((q-1)^{-1}[\varphi, \gamma]\big) \in \pi_{F_q}(\mathcal{I}_q) = \mathcal{I}$, which means that $\mathcal{I}$ is closed for the Poisson bracket. Thus (see (2.6)) K is coisotropic.

Similar arguments work when dealing with $\mathcal{C}_q$, $\mathfrak{I}_q$ or $\mathfrak{C}_q$. We shall only remark that working with $\mathcal{C}_q$ we end up with $\mathcal{C}\big(\widehat{K}\big) = \mathcal{C}(K) \leq_{\mathcal{P}} F[G]$, whence $\widehat{K}$ is coisotropic. $\square$

Since we would like to show also what happens in the non coisotropic case, we will consider, also, the weakest possible — naïve — version of quantization.

Definition 3.6. *Let $F_q[G]$ and $U_q(\mathfrak{g})$ be a QFA and a QUEA for G and $\mathfrak{g}$ and let*

$$\pi_{F_q}\colon\ F_q[G] \twoheadrightarrow F_q[G]\Big/(q-1)F_q[G] \ \cong\ F[G]$$
$$\pi_{U_q}\colon U_q(\mathfrak{g}) \twoheadrightarrow U_q(\mathfrak{g})\Big/(q-1)\,U_q(\mathfrak{g}) \ \cong\ U(\mathfrak{g})$$

be the specialization maps. Let $\mathcal{I}$, $\mathcal{C}$, $\mathfrak{I}$ and $\mathfrak{C}$ be the algebraic objects associated with the subgroup K of G (see 2.1). We call "weak quantization" of each of them any object $\mathcal{I}_q$, $\mathcal{C}_q$, $\mathfrak{I}_q$ or $\mathfrak{C}_q$ respectively, such that

$$\begin{array}{lccc}
(a) & \mathcal{I}_q \trianglelefteq_\ell \dot{\trianglelefteq}\, F_q[G] & , & \pi_{F_q}(\mathcal{I}_q) \,=\, \mathcal{I} \\
(b) & \mathcal{C}_q \leq^1 \dot{\trianglelefteq}_\ell\, F_q[G] & , & \pi_{F_q}(\mathcal{C}_q) \,=\, \mathcal{C} \\
(c) & \mathfrak{I}_q \trianglelefteq_\ell \dot{\trianglelefteq}\, U_q(\mathfrak{g}) & , & \pi_{U_q}(\mathfrak{I}_q) \,=\, \mathfrak{I} \\
(d) & \mathfrak{C}_q \leq^1 \dot{\trianglelefteq}_\ell\, U_q(\mathfrak{g}) & , & \pi_{U_q}(\mathfrak{C}_q) \,=\, \mathfrak{C}
\end{array} \tag{3.6}$$

It is obvious that strict or proper quantizations are weak. Let us remark that every subgroup of G is quantizable in the weak sense, since we may just consider e.g. $\mathcal{I}_q := \pi_{F_q}^{-1}(\mathcal{I})$ to be a quantization of $\mathcal{I}$. As naïf as it may seem, this remark will play a role in what follows.

Let us lastly remark how the real case should be treated.

Definition 3.7. *Let $(F_q[G], *)$ and $(U_q(\mathfrak{g}), *)$ be a real QFA and a real QUEA for G and $\mathfrak{g}$. Let $\mathcal{I}_q$, $\mathcal{C}_q$, $\mathfrak{I}_q$ and $\mathfrak{C}_q$ be subgroup quantizations (either strict, proper or weak). Then such quantizations are called real if*

$$(S(\mathcal{I}_q))^\star = \mathcal{I}_q\,, \quad \mathcal{C}_q^\star = \mathcal{C}_q\,, \quad (S(\mathfrak{I}_q))^\star = \mathfrak{I}_q\,, \quad \mathfrak{C}_q^\star = \mathfrak{C}_q \tag{3.7}$$

4 Quantum duality principle

Drinfeld's quantum duality principle (cf. [6], §7; see also [8] for a proof) has a stronger version (see [10]) best suited for *our* quantum groups (in the sense of Section 3.

Let H be any Hopf algebra in $\mathcal{HA}$ and let

$$I := Ker\Big(H \overset{\epsilon}{\twoheadrightarrow} \mathbb{C}\big[q,q^{-1}\big] \overset{ev_1}{\twoheadrightarrow} \mathbb{C}\Big) = Ker\Big(H \overset{ev_1}{\twoheadrightarrow} H/(q-1)H \overset{\bar{\epsilon}}{\twoheadrightarrow} \mathbb{C}\Big) \tag{4.1}$$

Then I is a Hopf ideal of H. We define

$$H^\vee \ :=\ \textstyle\sum_{n\geq 0}(q-1)^{-n}I^n \ =\ \bigcup_{n\geq 0}\big((q-1)^{-1}I\big)^n \ \ \Big(\subseteq \mathbb{C}(q)\otimes_{\mathbb{C}[q,q^{-1}]} H\Big) \tag{4.2}$$

Notice that, setting $J := Ker\Big(H \overset{\epsilon}{\twoheadrightarrow} \mathbb{C}\big[q,q^{-1}\big]\Big)$, one has $I = (q-1)\cdot 1_H + J$, so that

$$H^\vee \ =\ \textstyle\sum_{n\geq 0}(q-1)^{-n}J^n \ =\ \sum_{n\geq 0}\big((q-1)^{-1}J\big)^n \tag{4.3}$$

Consider, now, for every $n \in \mathbb{N}$ the iterated coproduct $\Delta^n\colon H \to H^{\otimes n}$ where

$$\Delta^0 := \epsilon \qquad \Delta^1 := \mathrm{id}_H \qquad \Delta^n := \big(\Delta\otimes \mathrm{id}_H^{\otimes(n-2)}\big)\circ\Delta^{n-1} \quad if\ n>2\ .$$

For any ordered subset $\Sigma = \{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$ with $i_1 < \dots < i_k$, define the morphism $j_\Sigma : H^{\otimes k} \longrightarrow H^{\otimes n}$ by

$$j_\Sigma(a_1 \otimes \dots \otimes a_k) := b_1 \otimes \dots \otimes b_n \text{ where } \begin{cases} b_i := 1 & \text{if} \quad i \notin \Sigma \\ b_{i_m} := a_m & \text{if} \quad 1 \leq m \leq k \end{cases}$$

then set $\Delta_\Sigma := j_\Sigma \circ \Delta^k$, $\Delta_\emptyset := \Delta^0$, and $\delta_\Sigma := \sum_{\Sigma' \subset \Sigma} (-1)^{n-|\Sigma'|} \Delta_{\Sigma'}$, $\delta_\emptyset := \epsilon$. By the inclusion-exclusion principle, the inverse formula $\Delta_\Sigma = \sum_{\Psi \subseteq \Sigma} \delta_\Psi$ holds. We shall use notation $\delta_0 := \delta_\emptyset$, $\delta_n := \delta_{\{1,2,\dots,n\}}$, and the key identity $\delta_n = (\mathrm{id_H} - \epsilon)^{\otimes n} \circ \Delta^n$, for all $n \in \mathbb{N}_+$. Given $H \in \mathcal{H}$, we define

$$H' := \left\{ a \in H \,\middle|\, \delta_n(a) \in (q-1)^n H^{\otimes n}, \ \forall\, n \in \mathbb{N} \right\} \quad \big(\subseteq H\big). \tag{4.4}$$

Theorem 4.1 (Global Quantum Duality Principle)**.** *(cf. [10]) For any $H \in \mathcal{HA}$ one has:*

(a) $H^\vee$ is a QUEA and H' is a QFA. Moreover the following inclusions hold:

$$H \subseteq \big(H^\vee\big)', \quad H \supseteq \big(H'\big)^\vee, \quad H^\vee = \big(\big(H^\vee\big)'\big)^\vee, \quad H' = \big(\big(H'\big)^\vee\big)' \tag{4.5}$$

(b) $H = \big(H^\vee\big)' \iff H$ is a QFA, and $H = \big(H'\big)^\vee \iff H$ is a QUEA;

(c) If G is a Poisson group with Lie bialgebra $\mathfrak{g}$, then

$$F_q[G]^\vee \Big/ (q-1) F_q[G]^\vee = U(\mathfrak{g}^*) \qquad U_q(\mathfrak{g})' / (q-1) U_q(\mathfrak{g})' = F[G^*]$$

where G^* is some connected Poisson group dual to G;

(d) Let $F_q[G]$ and $U_q(\mathfrak{g})$ be dual to each other w.r. to some perfect Hopf pairing. Then $F_q[G]^\vee$ and $U_q(\mathfrak{g})'$ are dual to each other w.r. to the same pairing.

A number of remarks are due, at this point:

1. The Poisson group G^* dual to G appearing in *(c)* of Theorem 4.1 does depend on $U_q(\mathfrak{g})$ which is given as a data. Different choices of $U_q(\mathfrak{g})$, though associated to the same Lie bialgebra $\mathfrak{g}$ may give rise to a different connected Poisson dual group G^*.

2. For all Hopf $\mathbb{C}(q)$–algebra $\mathbb{H}$ the existence of a $\mathbb{C}\big[q, q^{-1}\big]$-integral form H_f which is a QUEA at $q = 1$ is equivalent to the existence of a $\mathbb{C}\big[q, q^{-1}\big]$–integer form H_u which is a QFA at $q = 1$.

3. All claims above have obvious analogues in the real case.

4. If H is a Hopf algebra and $\Phi \subseteq \mathbb{N}$ is a finite subset, then ([14], Lemma 3.2)

$$\delta_\Phi(ab) = \sum_{\Lambda \cup Y = \Phi} \delta_\Lambda(a) \delta_\Phi(b) \qquad \forall a, b \in H \tag{4.6}$$

furthermore, if $\Phi \neq \emptyset$ we have

$$\delta_\Phi(ab - ba) = \sum_{\substack{\Lambda \cup Y = \Phi \\ \Lambda \cap Y \neq \emptyset}} \big(\delta_\Lambda(a)\, \delta_Y(b) - \delta_Y(b)\, \delta_\Lambda(a)\big) \qquad \forall a, b \in H \tag{4.7}$$

The above formulas will be used frequently in what follows

Having clarified the exact statement of quantum duality principle that we have in mind, let us extend it to objects of subgroup type as in Definition 3.6, i.e. to left coideal subalgebras and to left ideals and two-sided coideals — either in $F_q[G]$ or in $U_q(\mathfrak{g})$. This was already done in [4] where we only considered *local* (i.e. over $\mathbb{C}[[h]]$) quantizations. Let us remark that the quantum duality principle we have in mind not only exchanges the rôle of algebras of functions with that of universal enveloping algebras, but also exchanges the role of subgroups with that of homogeneous spaces. At the semiclassical level, the pair of dual objects is given by a coisotropic subgroup H and a Poisson quotient $G^* big/H^\perp$. When H is a Poisson subgroup, its orthogonal $H^\perp$ turns out to be normal in G^* and $G^*\big/H^\perp \cong H^*$ as a Poisson group, thus recovering the usual quantum duality principle. In particular, we will consider a process moving along the following draft:

$$
\begin{array}{lllllll}
(a) & \mathcal{I}\,\big(\subseteq F[G]\,\big) & \xrightarrow{(1)} & \mathcal{I}_q\,\big(\subseteq F_q[G]\,\big) & \xrightarrow{(2)} & \mathcal{I}_q^{\curlyvee}\,\big(\subseteq F_q[G]^\vee\,\big) & \xrightarrow{(3)}\ \mathcal{I}_1^{\curlyvee}\,\big(\subseteq U(\mathfrak{g}^*)\big) \\
(b) & \mathcal{C}\,\big(\subseteq F[G]\,\big) & \xrightarrow{(1)} & \mathcal{C}_q\,\big(\subseteq F_q[G]\,\big) & \xrightarrow{(2)} & \mathcal{C}_q^{\triangledown}\,\big(\subseteq F_q[G]^\vee\,\big) & \xrightarrow{(3)}\ \mathcal{C}_1^{\triangledown}\,\big(\subseteq U(\mathfrak{g}^*)\big) \\
(c) & \mathfrak{I}\,\big(\subseteq U(\mathfrak{g})\,\big) & \xrightarrow{(1)} & \mathfrak{I}_q\,\big(\subseteq U_q(\mathfrak{g})\,\big) & \xrightarrow{(2)} & \mathfrak{I}_q^{\,!}\,\big(\subseteq U_q(\mathfrak{g})'\,\big) & \xrightarrow{(3)}\ \mathfrak{I}_1^{\,!}\,\big(\subseteq F[G^*]\big) \\
(d) & \mathfrak{C}\,\big(\subseteq U(\mathfrak{g})\,\big) & \xrightarrow{(1)} & \mathfrak{C}_q\,\big(\subseteq U_q(\mathfrak{g})\,\big) & \xrightarrow{(2)} & \mathfrak{C}_q^{\,\Lsh}\,\big(\subseteq U_q(\mathfrak{g})'\,\big) & \xrightarrow{(3)}\ \mathfrak{C}_1^{\,\Lsh}\,\big(\subseteq F[G^*]\big)
\end{array}
$$

where arrows (1) are quantizations, arrows (3) are specializations at $q = 1$ and the definition of arrows (2) will be the core of what follows. It will turn out that:

1. each one of the right-hand-side objects above is one of the four algebraic objects which describe a closed connected subgroup of G^* : namely, the correspondence is

   (a) $\Longrightarrow$ *(c)* ,   *(b)* $\Longrightarrow$ *(d)* ,   *(c)* $\Longrightarrow$ *(a)* ,   *(d)* $\Longrightarrow$ *(b)* .

2. the four quantizations of subgroups of G^* so obtained are always *proper* — hence the subgroups of G^* associated to them are *coisotropic*.

3. if we begin with *strict* quantizations, and we start from a subgroup K, then the quantization of the unique coisotropic closed connected subgroup of G^* mentioned above is *strict* as well, and the subgroup itself is $K^\perp$ (cf. Definition 2.1), with some care in case *(b)*, i.e. if we start from $\mathcal{C}(K)$. This will partially generalize to *weak* quantizations, for which, starting from a subgroup K of G, the unique coisotropic closed connected subgroup of G^* obtained above is $K^{\langle\perp\rangle}$ (cf. Proposition 2.2).

Let us fix, in what follows, quantizations $U_q(\mathfrak{g})$ and $F_q[G]$ as in Section 3. Unless explicitely mentioned we will not assume that this is a double quantization. To simplify notations, let us set

$$
\begin{array}{ccc}
\mathbb{U}_q := \mathbb{U}_q(\mathfrak{g}) \quad , & U_q := U_q(\mathfrak{g}) \quad , & U_q{}' := U_q(\mathfrak{g})' \\
\mathbb{F}_q := \mathbb{F}_q[G] \quad , & F_q := F_q[G] \quad , & F_q{}^\vee := F_q[G]^\vee
\end{array}
$$

As mentioned in the first remark after Theorem 4.1, this implies that a specific connected Poisson dual G^* of G is selected (it depends on the choice of $U_q := U_Q(\mathfrak{g})$, not only on $\mathfrak{g}$ itself). Let us consider quantum subgroups $\mathcal{I}_q$, $\mathcal{C}_q$, $\mathfrak{I}_q$ and $\mathfrak{C}_q$ as defined in 3.6.

Definition 4.2. *Using notations as in (4.1) we define:*

(a) $\mathcal{I}_q^{\curlyvee} \; := \; \sum\limits_{n=1}^{\infty} (q-1)^{-n} \cdot I_{F_q}^{n-1} \cdot \mathcal{I}_q \; = \; \sum\limits_{n=1}^{\infty} (q-1)^{-n} \cdot J_{F_q}^{n-1} \cdot \mathcal{I}_q$

(b) $\mathcal{C}_q^{\triangledown} \; := \; \sum\limits_{n=0}^{\infty} (q-1)^{-n} \cdot \big(\mathcal{C}_q \cap I_{F_q}\big)^n \; = \; \sum\limits_{n=0}^{\infty} (q-1)^{-n} \cdot \big(\mathcal{C}_q \cap J_{F_q}\big)^n$

(c) $\mathfrak{I}_q^{\,!} \; := \; \Big\{ x \in \mathfrak{I}_q \;\Big|\; \delta_n(x) \in (q-1)^n \sum_{s=1}^n U_q^{\otimes(s-1)} \otimes \mathfrak{I}_q \otimes U_q^{\otimes(n-s)}, \; \forall \, n \in \mathbb{N}_+ \Big\}$

(d) $\mathfrak{C}_q^{\,\Lsh} \; := \; \Big\{ x \in \mathfrak{C}_q \;\Big|\; \delta_n(x) \in (q-1)^n \, U_q^{\otimes(n-1)} \otimes \mathfrak{C}_q \,, \; \forall \, n \in \mathbb{N}_+ \Big\}$

Let us remark that the following inclusions hold directly by definitions:

$$(i) \;\; \mathcal{I}_q^{\curlyvee} \supseteq \mathcal{I}_q \,, \qquad (ii) \;\; \mathcal{C}_q^{\triangledown} \supseteq \mathcal{C}_q \,, \qquad (iii) \;\; \mathfrak{I}_q^{\,!} \subseteq \mathfrak{I}_q \,, \qquad (iv) \;\; \mathfrak{C}_q^{\,\Lsh} \subseteq \mathfrak{C}_q \,. \tag{4.8}$$

5 Duality maps

In the present section we will prove properties of the four Drinfeld–type maps defined in the previous section, namely the maps $\mathcal{I}_q \mapsto \mathcal{I}_q^{\curlyvee}$, $\mathcal{C}_q \mapsto \mathcal{C}_q^{\triangledown}$, $\mathfrak{I}_q \mapsto \mathfrak{I}_q^{\,!}$ and $\mathfrak{C}_q \mapsto \mathfrak{C}_q^{\,\Lsh}$. Let us recall that such maps do not change, as we will see, the algebraic properties of subobjects, but interchanges quantized function algebra with quantum enveloping algebra and therefore quantizations of coisotropic subgroups will be sent to quantizations of (embedebbable) homogeneous spaces — of the dual quantum group — and viceversa.

Let us start by considering the map $\mathcal{I}_q \mapsto \mathcal{I}_q^{\curlyvee}$.

Proposition 5.1. *Let $\mathcal{I}_q = \mathcal{I}_q(K)$ be a left coideal and two-sided coideal in $F_q[G]$, that is a weak quantization (of type $\mathcal{I}$) of some subgroup K of G. Then*

1. *$\mathcal{I}_q^{\curlyvee}$ is a left ideal and two-sided coideal in $F_q[G]^{\vee}$;*
2. *if $\mathcal{I}_q$ is strict, then $\mathcal{I}_q^{\curlyvee}$ is strict too, i.e. $\mathcal{I}_q^{\curlyvee} \bigcap \, (q-1) \, F_q[G]^{\vee} \; = \; (q-1) \, \mathcal{I}_q^{\curlyvee}$;*
3. *there exists a coisotropic subgroup L of G^* such that $\mathcal{I}_q(K)^{\curlyvee} = \mathfrak{I}_q(L)$: namely, $\mathcal{I}_q(K)^{\curlyvee}$ is a proper quantization, of type $\mathfrak{I}$, of some coisotropic subgroup L of G^* ;*
4. *in the real case, i.e. if the quantization $\mathcal{I}_q$ is a real one, $\mathcal{I}_q^{\curlyvee}$ is real too, i.e. $\big(S\big(\mathcal{I}_q^{\curlyvee}\big)\big)^* = \mathcal{I}_q^{\curlyvee}$. Therefore claims (1–3) still hold in the framework of real quantum subgroups.*

Proof. (1) Consider that $\mathcal{I}_q^{\curlyvee}$ is the left ideal of $F_q^{\vee}$ generated by $(q-1)^{-1} \mathcal{I}_q$; therefore, in order to prove $\mathcal{I}_q^{\curlyvee} \,\dot{\trianglelefteq}\, F_q^{\vee}$ it is enough to show that $\Delta\big((q-1)^{-1} \mathcal{I}_q\big) \subseteq F_q^{\vee} \otimes \mathcal{I}_q^{\curlyvee} + \mathcal{I}_q^{\curlyvee} \otimes F_q^{\vee}$. Since $\mathcal{I}_q$ is a coideal of F_q, we have

$$\Delta\big((q-1)^{-1} \mathcal{I}_q\big) \subseteq F_q \otimes (q-1)^{-1} \mathcal{I}_q + (q-1)^{-1} \mathcal{I}_q \otimes F_q \subseteq F_q^{\vee} \otimes \mathcal{I}_q^{\curlyvee} + \mathcal{I}_q^{\curlyvee} \otimes F_q^{\vee} \tag{5.1}$$

whence ${\mathcal{I}_q}^{\curlyvee} \dot{\trianglelefteq} {F_q}^{\vee}$ follows, and the first claim is proved.

(2) Assume $\mathcal{I}_q$ to be a *strict* quantization, so that $\mathcal{I}_q \bigcap (q-1)\, F_q \, = \, (q-1)\, \mathcal{I}_q$. Let $J := \mathit{Ker}\big(\epsilon : \, F_q \longrightarrow \mathbb{C}\big[q,q^{-1}\big]\big)$. Then

$$J \mod (q-1) F_q = \mathit{Ker}\,(\epsilon)\big|_{F[G]} \; = \; \mathfrak{m}_e$$

and $\mathfrak{m}_e \Big/ {\mathfrak{m}_e}^2 = \mathfrak{g}^*$, the cotangent Lie bialgebra of G . Let $\{y_1, \dots, y_n\}$ be a subset of $\mathfrak{m}_e$ whose image in the local ring of G at the identity e is a local system of parameters, and pull it back to a subset $\{j_1, \dots, j_n\}$ of J. Let $\widehat{F}_q$ be the J–adic completion of F_q. From [10], Lemma 4.1, we know that the set of ordered monomials $\big\{ j^{\,\underline{e}} \;\big|\; \underline{e} \in \mathbb{N}^n \big\}$ (where hereafter $j^{\,\underline{e}} := \prod_{s=1}^n j_s^{\underline{e}(i)}$, for all $\underline{e} \in \mathbb{N}^n$) is a $\mathbb{C}\big[q,q^{-1}\big]$–pseudobasis of $\widehat{F}_q$, which means that each element of $\widehat{F}_q$ has a unique expansion as a formal infinite linear combination of the $j^{\,\underline{e}}$'s. In a similar way, the $(q-1)$–adic completion of ${F_q}^{\vee}$ admits $\big\{ (q-1)^{-|\underline{e}|} j^{\,\underline{e}} \;\big|\; \underline{e} \in \mathbb{N}^n \big\}$ as a $\mathbb{C}\big[q,q^{-1}\big]$–pseudobasis, where $|\underline{e}| := \sum_{i=1}^n \underline{e}(i)$.

For our purposes we need a special choice of the set $\{j_1, \dots, j_n\}$ adapted to the smooth subvariety K of G. By general theory we can choose $\{y_1, \dots, y_n\}$ so that $y_1, \, \dots, \, y_k \in \mathfrak{m}_e$ and $y_{k+1}, \dots, y_n \in \mathcal{I}(K)$, where $k = \mathit{dim}(K)$. We can also choose the lift $\{j_1, \dots, j_n\}$ of $\{y_1, \dots, y_n\}$ inside J so that j_s is a lift of y_s, for all $s = 1, \dots, k$, and $j_{k+1}, \dots, j_n \in \mathcal{I}_q$. With these assumptions, it's easy to see that

$$\varphi \in {\mathcal{I}_q}^{\curlyvee} \bigcap (q-1)\, {F_q}^{\vee} \;\; \Longrightarrow \;\; (q-1)^n \, \varphi \in \big(J^{n-1} \cdot \mathcal{I}_q\big) \bigcap (q-1)\, J^n$$

for some $n \in \mathbb{N}$, which in turn yields $(q-1)^n \varphi \in J^{n-1} \cdot \big(\mathcal{I}_q \bigcap (q-1)\, J\big)$. Since

$$\mathcal{I}_q \bigcap (q-1)\, J \; \subseteq \; \mathcal{I}_q \bigcap (q-1)\, F_q \; = \; (q-1)\, \mathcal{I}_q$$

we conclude that $(q-1)^n \, \varphi \in (q-1)\, J^{n-1} \cdot \mathcal{I}_q$, whence $\varphi \in (q-1)\, {\mathcal{I}_q}^{\curlyvee}$. The converse inclusion ${\mathcal{I}_q}^{\curlyvee} \bigcap (q-1)\, {F_q}^{\vee} \supseteq (q-1)\, {\mathcal{I}_q}^{\curlyvee}$ is obvious, hence claim *(2)* is proved.

(3) It is an obvious statement that ${\mathcal{I}_q}^{\curlyvee}$ is a weak quantization of its image $\pi_{{F_q}^{\vee}}\big({\mathcal{I}_q}^{\curlyvee}\big)$: in particular, $\pi_{{F_q}^{\vee}}\big({\mathcal{I}_q}^{\curlyvee}\big) \trianglelefteq_{\ell} \dot{\trianglelefteq} \; \pi_{{F_q}^{\vee}}\big({F_q}^{\vee}\big) \; = \; U\big(\mathfrak{g}^*\big)$ implies that $\pi_{{F_q}^{\vee}}\big({\mathcal{I}_q}^{\curlyvee}\big) \, = \, \mathfrak{I}(L)$ for some subgroup L of G^*. Thus ${\mathcal{I}_q}^{\curlyvee}$ is a weak quantization, to be called $\mathfrak{I}_q(L)$, of $\mathfrak{I}(L)$, and it is even strict if $\mathcal{I}_q$ itself is strict, as we've just seen. Now we show that such quantization $\mathfrak{I}_q(L)$ turns out to be always *proper*.

In fact, (5.1) implies $\nabla\big((q-1)^{-1} \mathcal{I}_q\big) \subseteq (q-1)^{-1} \big(F_q \wedge \mathcal{I}_q\big)$. On the other hand $F_q \wedge \mathcal{I}_q \subseteq J \wedge \mathcal{I}_q \subseteq (q-1)^2 \, {F_q}^{\vee} \wedge {\mathcal{I}_q}^{\curlyvee}$, thus, finally, $\nabla\big({\mathcal{I}_q}^{\curlyvee}\big) \in (q-1)\, {F_q}^{\vee} \wedge {\mathcal{I}_q}^{\curlyvee}$, which means that ${\mathcal{I}_q}^{\curlyvee}$ is proper and *(3)* holds.

(4) This is an obvious consequence of definitions. □

Remark 5.2. In functorial language we may say that the map $\mathcal{I}_q \mapsto {\mathcal{I}_q}^{\curlyvee}$ establishes a functor between quantizations of coisotropic subgroups of G and quantizations of (embeddable) homogeneous spaces of G^*, moving from a global to a local description, sending each type of quantization in a proper one and preserving strictness. Indeed, we should make precise what are the "arrows" in our categories of "quantum subgroups" or "quantum homogeneous spaces", and how the functor acts on these: we leave these details to the interested reader.

Let us move on to properties of the map $\mathcal{C}_q \mapsto \mathcal{C}_q^{\triangledown}$.

Proposition 5.3. *Let $\mathcal{C}_q = \mathcal{C}_q(K)$ be a left coideal subalgebra in $F_q[G]$. Then*

1. *$\mathcal{C}_q^{\triangledown}$ is a left coideal subalgebra in $F_q[G]^{\vee}$;*
2. *if $\mathcal{C}_q$ is strict, then $\mathcal{C}_q^{\triangledown}$ is strict too, i.e. $\mathcal{C}_q^{\triangledown} \bigcap (q-1)\, F_q[G]^{\vee} \,=\, (q-1)\, \mathcal{C}_q^{\triangledown}$.*
3. *there exists a coisotropic subgroup L of G^* such that $\mathcal{C}_q(K)^{\triangledown} = \mathfrak{C}_q(L)$: namely, $\mathcal{C}_q(K)^{\triangledown}$ is a proper quantization, of type $\mathfrak{C}$, of some coisotropic subgroup L of G^* ;*
4. *in the real case, i.e. if the quantization $\mathcal{C}_q$ is a real one, $\mathcal{C}_q(K)^{\triangledown}$ is real too, i.e. $\big(\mathcal{C}_q^{\triangledown}\big)^* = \mathcal{C}_q^{\triangledown}$. Therefore claims (1–3) still hold in the framework of real quantum subgroups.*

Proof. The proof uses essentially the same arguments as the previous one.

(1) By the very definitions $\mathcal{C}_q^{\triangledown} \leq^1 F_q^{\vee} := F_q[G]^{\vee}$. More precisely, $\mathcal{C}_q^{\triangledown}$ is (by construction) the unital $\mathbb{C}\big[q,q^{-1}\big]$–subalgebra of $F_q^{\vee}$ generated by $(q-1)^{-1}\,(\mathcal{C}_q)^+$, where $(\mathcal{C}_q)^+ := \mathcal{C}_q \bigcap J$. So to get $\mathcal{C}_q^{\triangledown} \dot{\trianglelefteq}_\ell\, F_q^{\vee}$ we must only prove $\Delta\big((q-1)^{-1}(\mathcal{C}_q)^+\big) \subseteq F_q^{\vee} \otimes \mathcal{C}_q^{\triangledown}$. But $\mathcal{C}_q \dot{\trianglelefteq}_\ell\, F_q$, so:

$$\Delta\big((q-1)^{-1}(\mathcal{C}_q)^+\big) \;\subseteq\; F_q \otimes (q-1)^{-1}(\mathcal{C}_q)^+ \;\subseteq\; F_q^{\vee} \otimes \mathcal{C}_q^{\triangledown} \tag{5.2}$$

therefore $\mathcal{C}_q^{\triangledown} \dot{\trianglelefteq}_\ell\, F_q^{\vee}$, and claim *(1)* is proved.

(2) Now suppose $\mathcal{C}_q$ to be a *strict* quantization, i.e. $\mathcal{C}_q \bigcap (q-1)\, F_q \,=\, (q-1)\, \mathcal{C}_q$. We need an explicit description of $F_q^{\vee}$ and of $\mathcal{C}_q^{\triangledown}$. This goes along the same lines followed to describe $\mathcal{I}_q^{\curlyvee}$ in the proof of Proposition 4.2: but now the choice of the subset $\{j_1, \dots, j_n\}$ of J is different.

First, since $\mathcal{C}(K) = \mathcal{C}\big(\widehat{K}\big)$ we can assume that $K = \widehat{K}$, i.e. K is observable. Then we can choose $\big\{j_1, \dots, j_n\big\}$ so that $j_{k+1}, \dots, j_n \in J \bigcap \mathcal{C}_q = \mathcal{C}_q^{\,+}$ (where again $k = dim\,(K)$) and, letting $y_s := j_s \mod (q-1)\, F_q$, the set $\big\{y_1, \dots, y_n\big\}$ yields a local system of parameters at $e \in G$ (in the localized ring), as before; now in addition we have $y_{k+1}, \dots, y_n \in \mathfrak{m}_e \bigcap \mathcal{C}(K) =: \mathcal{C}(K)^+$. With these assumptions, the $(q-1)$–adic completion of $F_q^{\vee}$ admits $\big\{ (q-1)^{-|\underline{e}|} j^{\,\underline{e}} \,\big|\, \underline{e} \in \mathbb{N}^{\,n} \big\}$ as a $\mathbb{C}\big[q,q^{-1}\big]$–pseudobasis, like before, but in addition the same analysis can be done for the $(q-1)$–adic completion of $\mathcal{C}_q^{\triangledown}$ (just because $\mathcal{C}_q$ is *strict*), which then has $\mathbb{C}\big[q,q^{-1}\big]$–pseudobasis $\big\{ \prod_{s=k+1}^n j_s^{\,e_s} \,\big|\, (e_{k+1}, \dots, e_n) \in \mathbb{N}^{\,n-k} \big\}$. From these description of the completions, and comparing the former with $F_q^{\vee}$ and $\mathcal{C}_q$, we easily see that $\mathcal{C}_q^{\triangledown} \bigcap (q-1)\, F_q^{\vee} \subseteq (q-1)\, \mathcal{C}_q^{\triangledown}$. The converse is trivial, hence claim *(1)* is proved.

(3) It follows directly from *(1)* that $\mathcal{C}_q^{\triangledown}$ is a weak quantization of its image $\pi_{F_q^{\vee}}\big(\mathcal{C}_q^{\triangledown}\big)$: in particular, $\pi_{F_q^{\vee}}\big(\mathcal{C}_q^{\triangledown}\big) \leq^1 \dot{\trianglelefteq}_\ell\, \pi_{F_q^{\vee}}\big(F_q^{\,\vee}\big) \,=\, U\big(\mathfrak{g}^*\big)$ means that $\pi_{F_q^{\vee}}\big(\mathcal{C}_q^{\triangledown}\big) = \mathfrak{C}(L)$ for some subgroup L of G^*. Thus $\mathcal{C}_q^{\triangledown}$ is a weak quantization — to be called $\mathfrak{C}_q(L)$ — of $\mathfrak{C}(L)$, and it is even strict if $\mathcal{C}_q$ itself is strict, by claim *(1).* Now in addition we show that, in any case, such a quantization $\mathfrak{C}_q(L)$ is always *proper*.

From (5.2) we have

$$\nabla\big((q-1)^{-1}\,(\mathcal{C}_q)^+\big) \;\subseteq\; (q-1)^{-1} J \wedge (\mathcal{C}_q)^+ \subseteq\; (q-1)^{-1+2}\, F_q^{\,\vee} \wedge \mathcal{C}_q^{\triangledown} \;=\; (q-1)\, F_q^{\,\vee} \wedge \mathcal{C}_q^{\triangledown}$$

which implies exactly that $\mathcal{C}_q^{\,\triangledown}$ — which by definition is the unital subalgebra generated by $(q-1)^{-1}\,(\mathcal{C}_q)^+$ — is proper.

(4) This follows directly from definitions and from $\mathcal{C}_q^{\,*} = \mathcal{C}_q$, which holds by assumption. ☐

Remark 5.4. In functorial language we may say that the map $\mathcal{C}_q \mapsto \mathcal{C}_q^{\,\triangledown}$ establishes a functor between quantized homogeneous spaces of G and quantizations of coisotropic subgroups of G^*, moving from a global to a local description, sending each type of quantization in a proper one and preserving strictness. Again, to be precise, several details need to be fixed, and are left to the reader.

The third step copes with the map $\mathfrak{I}_q \mapsto \mathfrak{I}_q^{\,!}$.

Proposition 5.5. *Let* $\mathfrak{I}_q = \mathfrak{I}_q(K)$ *be a left ideal and two-sided coideal in* $U_q(\mathfrak{g})$ *, weak quantization (of type* $\mathfrak{I}$*) of some coisotropic subgroup* K *of* G *. Then:*

1. $\mathfrak{I}_q^{\,!}$ *is a left ideal and two-sided coideal in* $U_q(\mathfrak{g})'$*;*
2. *if* $\mathfrak{I}_q$ *is strict, then* $\mathfrak{I}_q^{\,!}$ *is strict too, i.e.* $\mathfrak{I}_q^{\,!} \bigcap (q-1)\, U_q(\mathfrak{g})' \, = \, (q-1)\, \mathfrak{I}_q^{\,!}$ *;*
3. *there exists a coisotropic subgroup* L *in* G^* *such that* $\mathfrak{I}_q(K)^! = \mathcal{I}_q(L)$ *: namely,* $\mathfrak{I}_q(K)^!$ *is a proper quantization, of type* $\mathcal{I}$ *, of some coisotropic subgroup* L *of* G^* *;*
4. *in the real case, i.e. if the quantization* $\mathfrak{I}_q$ *is a real one,* $\mathfrak{I}_q^{\,!}$ *is real too, i.e.* $\big(S\big(\mathfrak{I}_q^{\,!}\big)\big)^* = \mathfrak{I}_q^{\,!}$*. Therefore claims (1–3) still hold in the framework of real quantum subgroups.*

Proof. (1) Let $a \in {U_q}'$ and $b \in \mathfrak{I}_q^{\,!}$: by definition of $\mathfrak{I}_q^{\,!}$, from $\mathfrak{I}_q \trianglelefteq_\ell U_q$ and from (4.6) we get

$$\delta_n(ab) \, \in \, (q-1)^n \sum_{s=1}^{n} {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)}$$

so $a\,b \in \mathfrak{I}_q^{\,!}$, thus $\mathfrak{I}_q^{\,!} \trianglelefteq_\ell {U_q}'$.

As to the coideal property, it is proven resorting to $(q-1)$–adic completions, arguing as in the proof of Proposition 3.5 in [10], and basing on the fact that $\mathfrak{I}_q \dot{\trianglelefteq}\, U_q$. Details are left to the reader.

(2) Assume now $\mathfrak{I}_q$ to be *strict*. The inclusion

$$\mathfrak{I}_q^{\,!} \bigcap (q-1)\, U_q(\mathfrak{g})' \, \supseteq \, (q-1)\, \mathfrak{I}_q^{\,!}$$

is trivially true, and we must prove the converse. Let $\eta \in \mathfrak{I}_q^{\,!} \bigcap (q-1)\, U_q(\mathfrak{g})'$. We have

$$\delta_n(\eta) \, \in \, (q-1)^n \left(\left(\textstyle\sum_{s=1}^{n} {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)} \right) \bigcap\, (q-1)\, {U_q}^{\otimes n} \right)$$

for all $n \in \mathbb{N}_+$. But then our assumption gives

$$\bigg(\sum_{s=1}^n {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)} \bigg) \bigcap \, (q-1) \, {U_q}^{\otimes n} =$$
$$= \sum_{s=1}^n {U_q}^{\otimes (s-1)} \otimes \Big(\mathfrak{I}_q \bigcap (q-1) \, U_q \Big) \otimes {U_q}^{\otimes (n-s)} \, = \, (q-1)^{n+1} \sum_{s=1}^n {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)}$$

which, in turn, means $\eta \in (q-1) \, {\mathfrak{I}_q}^{\!!}$. Thus ${\mathfrak{I}_q}^{\!!} \bigcap (q-1) \, U_q(\mathfrak{g})' \subseteq (q-1) \, {\mathfrak{I}_q}^{\!!}$, as expected.

(3) Claim *(1)* implies that ${\mathfrak{I}_q}^{\!!}$ is a weak quantization of its image, therefore there exists a subgroup L of G^* such that $\pi_{U_q'}\big({\mathfrak{I}_q}^{\!!}\big) = \mathcal{I}(L)$. This quantization is even strict if $\mathfrak{I}_q$ itself is strict, by the previous. Now we show that this quantization $\mathcal{I}_q(L)$ is always *proper* — hence the subgroup L is coisotropic, by Lemma 3.5.

Recall that, by definition, $\mathcal{I}_q(L)$ is proper if and only if $[x,y] \in (q-1) \, {\mathfrak{I}_q}^{\!!}$ for all $x, y \in {\mathfrak{I}_q}^{\!!}$. From definitions we have

$$[x,y] \in (q-1) \, {\mathfrak{I}_q}^{\!!} \iff \delta_n\big([x,y]\big) \in (q-1)^{n+1} \textstyle\sum_{s=1}^n {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)} \quad \forall \, n \in \mathbb{N}$$

Then by formula (4.7) we have (for all $n \in \mathbb{N}$)

$$\delta_n\big([x,y]\big) \; = \; \textstyle\sum_{\substack{\Lambda \cup Y = \{1,\dots,n\} \\ \Lambda \cap Y \neq \emptyset}} \big(\, \delta_\Lambda(x) \, \delta_Y(y) \, - \, \delta_Y(y) \, \delta_\Lambda(x) \, \big) \tag{5.3}$$

while (with notation of §4)

$$\delta_\Lambda(x) \in (q-1)^{|\Lambda|} \cdot j_\Lambda \left(\textstyle\sum_{s=1}^{|\Lambda|} {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (|\Lambda|-s)} \right) \, ,$$
$$\delta_Y(y) \in (q-1)^{|Y|} \cdot j_Y \left(\textstyle\sum_{s=1}^{|Y|} {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (|Y|-s)} \right) \, ;$$

since $\Lambda \cup Y = \{1,\dots,n\}$ and $\Lambda \cap Y \neq \emptyset$ we have $|\Lambda| + |Y| \geq n+1$; moreover, for each index $i \in \{1,\dots,n\}$ we have $i \in \Lambda$ (and otherwise *Im* (j_Λ) has 1 in the i–th spot) or $i \in Y$ (with the like remark on *Im* (j_Y) if not). As $\mathfrak{I}_q$ is a left ideal of U_q, we conclude

$$\begin{aligned} \delta_\Lambda(x) \cdot \delta_Y(y) \, , \; \delta_Y(y) \cdot \delta_\Lambda(x) \; \in \; & (q-1)^{|\Lambda|+|Y|} \textstyle\sum_{s=1}^n {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)} \\ & \subseteq (q-1)^{n+1} \textstyle\sum_{s=1}^n {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)} \end{aligned}$$

so that (5.3) gives $\delta_n\big([x,y]\big) \in (q-1)^{n+1} \sum_{s=1}^n {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)}$, as expected.

(4) In the real case, $\big(S\big({\mathfrak{I}_q}^{\!!}\big) \big)^* = {\mathfrak{I}_q}^{\!!}$ follows at once from definitions and from $\big(S(\mathfrak{I}_q) \big)^* = \mathfrak{I}_q$. $\square$

Remark 5.6. In functorial language we may say that the map $\mathfrak{I}_q \mapsto {\mathfrak{I}_q}^{\!!}$ establishes a functor between quantized homogeneous spaces of G and quantizations of coisotropic subgroups of G^*, moving from a local to a global description, sending each type of quantization in a proper one and preserving strictness. Once more, details are left to the interested reader.

The fourth and last step is devoted to the map $\mathfrak{C}_q \mapsto {\mathfrak{C}_q}^{\Lsh}$.

Proposition 5.7. *Let $\mathfrak{C}_q = \mathfrak{C}_q(K)$ be a subalgebra and left coideal in $U_q(\mathfrak{g})$, weak quantization (of type $\mathfrak{C}$) of some subgroup K of G. Then:*

1. *$\mathfrak{C}_q^{\Lsh}$ is a subalgebra and left coideal in $U_q(\mathfrak{g})'$;*
2. *if $\mathfrak{C}_q$ is strict, then $\mathfrak{I}_q^{\,!}$ is strict too, i.e. $\mathfrak{C}_q^{\Lsh} \bigcap (q-1)\, U_q(\mathfrak{g})' \, = \, (q-1)\, \mathfrak{C}_q^{\Lsh}$;*
3. *there exists a coisotropic subgoup L in G^* such that $\mathfrak{C}_q(K)^{\Lsh} = \mathcal{C}_q(L)$: namely, $\mathfrak{C}_q(K)^{\Lsh}$ is a proper quantization, of type $\mathcal{C}$, of some coisotropic subgroup L of G^* ;*
4. *in the real case, i.e. if the quantization $\mathfrak{C}_q$ is a real one, $\mathfrak{C}_q(K)^{\Lsh}$ is real too, i.e. $\big(\mathfrak{C}_q^{\Lsh}\big)^* = \mathfrak{C}_q^{\Lsh}$. Therefore claims (1–3) still hold in the framework of real quantum subgroups.*

Proof. The whole proof is very similar to that of Proposition 5.5.

(1) By definitions, $1 \in \mathfrak{C}_q$ and $\delta_n(1) = 0$ for all $n \in \mathbb{N}$, so $1 \in \mathfrak{C}_q^{\Lsh}$. Let $x, y \in \mathfrak{C}_q^{\Lsh}$ and $n \in \mathbb{N}$; by (4.6) we have $\delta_n(xy) = \sum_{\Lambda \cup Y = \{1,\dots,n\}} \delta_\Lambda(x)\, \delta_Y(y)$. Each of the factors $\delta_\Lambda(x)$ belongs to a module $(q-1)^{|\Lambda|}\, {U_q}^{\otimes(|\Lambda|-1)} \otimes X$ where the last tensor factor is either $X = \mathfrak{C}_q$ (if $n \in \Lambda$) or $X = \{1\} \subset \mathfrak{C}_q$ (if $n \not\in \Lambda$), and similarly for $\delta_Y(y)$; in addition $\Lambda \cup Y = \{1,\dots,n\}$ implies $|\Lambda| + |Y| \geq n$, and summing up $\delta_n(xy) \in (q-1)^n {U_q}^{\otimes(n-1)} \otimes \mathfrak{C}_q$, whence $x\,y \in \mathfrak{C}_q^{\Lsh}$. Thus $\mathfrak{C}_q^{\Lsh}$ is a subalgebra of ${U_q}'$.

In order to prove that $\mathfrak{C}_q^{\Lsh}$ is a left coideal in ${U_q}'$, one can again resort to $(q-1)$–adic completions, with exactly the same arguments as in the proof of Proposition 3.5 in [4], starting from the fact that $\mathfrak{C}_q \,\dot{\trianglelefteq}_\ell\, U_q$. Details are left to the reader.

(2) Assume, now, that $\mathfrak{C}_q$ is a *strict* quantization, i.e. $\mathfrak{C}_q \bigcap (q-1)\, F_q \, = (q-1)\, \mathfrak{C}_q$. Then clearly $\mathfrak{C}_q^{\Lsh} \bigcap (q-1)\, U_q(\mathfrak{g})' \supseteq (q-1)\, \mathfrak{C}_q^{\Lsh}$, and we must prove the converse inclusion. Let $\kappa \in \mathfrak{C}_q^{\Lsh} \bigcap (q-1)\, U_q(\mathfrak{g})'$. Then:

$$\delta_n(\kappa) \,\in\, (q-1)^n \Big(\Big({U_q}^{\otimes(n-1)} \otimes \mathfrak{C}_q \Big) \bigcap (q-1)\, {U_q}^{\otimes n} \Big) \,=$$
$$= \, (q-1)^n \Big({U_q}^{\otimes(n-1)} \otimes \big(\mathfrak{C}_q \bigcap (q-1)\, U_q \big) \Big) \,=\, (q-1)^{n+1} \cdot {U_q}^{\otimes(n-1)} \otimes \mathfrak{C}_q$$

which means $\kappa \in (q-1)\, \mathfrak{C}_q^{\Lsh}$. Therefore $\mathfrak{C}_q^{\Lsh} \bigcap (q-1)\, U_q(\mathfrak{g})' \subseteq (q-1)\, \mathfrak{C}_q^{\Lsh}$, as claimed.

(3) The above algebraic properties show that $\mathfrak{C}_q^{\Lsh}$ is a weak quantization of its image $\pi_{{U_q}'}\big(\mathfrak{C}_q^{\Lsh}\big)$; thus there exists a coisotropic subgroup L of G^* such that: $\pi_{{U_q}'}\big(\mathfrak{C}_q^{\Lsh}\big) = \mathcal{C}(L)$. Thus $\mathfrak{I}_q^{\,!}$ is a weak quantization — to be called $\mathcal{I}_q(L)$ — of $\mathcal{I}(L)$, and it is even strict if $\mathfrak{I}_q$ itself is strict, by the previous. Now we show first that this quantization $\mathcal{I}_q(L)$ is always *proper* — hence the subgroup L is coisotropic, by Lemma 3.5.

Proving that $\mathcal{I}_q(L)$ is proper amounts to show that $[x,y] \in (q-1)\, \mathfrak{C}_q^{\Lsh}$ for all $x, y \in \mathfrak{C}_q^{\Lsh}$. By definition we have

$$[x,y] \,\in\, (q-1)\, \mathfrak{C}_q^{\Lsh} \iff \delta_n\big([x,y]\big) \,\in\, (q-1)^{n+1} {U_q}^{\otimes(n-1)} \otimes \mathfrak{C}_q \quad \forall\, n \in \mathbb{N}$$

and formula (4.7) gives, for all $n \in \mathbb{N}$,

$$\delta_n\big([x,y]\big) \,=\, \textstyle\sum_{\substack{\Lambda \cup Y = \{1,\dots,n\} \\ \Lambda \cap Y \neq \emptyset}} \big(\delta_\Lambda(x)\, \delta_Y(y) \,-\, \delta_Y(y)\, \delta_\Lambda(x) \big) \tag{5.4}$$

while

$$\delta_\Lambda(x) \,\in\, (q-1)^{|\Lambda|}\, j_\Lambda\Big({U_q}^{\otimes(|\Lambda|-1)} \otimes \mathfrak{C}_q\Big) \;, \quad \delta_Y(y) \,\in\, (q-1)^{|Y|}\, j_Y\Big({U_q}^{\otimes(|Y|-1)} \otimes \mathfrak{C}_q\Big) \;.$$

Now, $\Lambda \cup Y = \{1, \dots, n\}$ and $\Lambda \cap Y \neq \emptyset$ give $|\Lambda| + |Y| \geq n+1$, and since $\mathfrak{C}_q$ is a subalgebra of U_q we get

$$\delta_\Lambda(x)\,\delta_Y(y) \,,\; \delta_Y(y)\,\delta_\Lambda(x) \,\in\, (q-1)^{|\Lambda|+|Y|}\,{U_q}^{\otimes(n-1)} \otimes \mathfrak{C}_q \subseteq (q-1)^{n+1}{U_q}^{\otimes(n-1)} \otimes \mathfrak{C}_q$$

so that (5.4) yields

$$\delta_n\big([x,y]\big) \,\in\, (q-1)^{n+1}{U_q}^{\otimes(n-1)} \otimes \mathfrak{C}_q$$

thus $[x,y] \in (q-1)\,\mathfrak{C}_q^{\,\Lsh}$.

(4) In the real case $(\mathfrak{C}_q)^* = \mathfrak{C}_q$: this and the very definitions imply the claim. $\square$

Remark 5.8. In functorial language we may say that the map $\mathfrak{C}_q \mapsto \mathfrak{C}_q^{\,\Lsh}$ establishes a functor between quantization of coisotropic subgroups of G and quantizations of Poisson homogeneoeous spaces of G^*, moving from a local to a global description, sending each type of quantization in a proper one and preserving strictness. We leave to the interested reader all details which still need to be fixed.

We now move to connectedness properties of the coisotropic subgroup L identified in Propositions 5.5 and 5.7.

Proposition 5.9.

1. *Let $\mathfrak{I}_q(K)$ be a strict quantization (of type $\mathfrak{I}$) of a (coisotropic) subgroup K in G. Then the subgroup L of G^* such that $\mathfrak{I}_q(K)^! = \mathcal{I}_q(L)$ is connected.*

2. *Let $\mathfrak{C}_q(K)$ be a strict quantization of type $\mathfrak{C}$ of a (coisotropic) subgroup K of G. Then the subgroup L of G^* such that $\mathfrak{C}_q(K)^! = \mathcal{C}_q(L)$ is connected.*

Proof. (1) Saying that the (closed) subgroup L is connected is equivalent to saying that its function algebra $F[L] = F\big[G^*\big]\Big/\mathcal{I}(L)$ has no non-trivial idempotents. Note that, since $F\big[G^*\big]$ is the specialization of U_q' at $q=1$ and $\mathcal{I}(L)$ is the similar specialization of $\mathfrak{I}_q^{\,!}$, the quotient $F[L] = F\big[G^*\big]\Big/\mathcal{I}(L)$ is canonically isomorphic to the specialization at $q=1$ of $U_q'\Big/\,\mathfrak{I}_q^{\,!}$. Let $\overline{a}$ be an idempotent in $F[L]$: if we take any lift of it in $U_q'\Big/\,\mathfrak{I}_q^{\,!}$, i.e. any $a \in U_q'\Big/\,\mathfrak{I}_q^{\,!}$ such that $\overline{a} = a \mod (q-1)\,U_q'\Big/\,\mathfrak{I}_q^{\,!}$. We must prove:

$$a^2 \,\equiv\, a \mod (q-1)\,U_q'\Big/\,\mathfrak{I}_q^{\,!} \quad\Longrightarrow\quad a \mod (q-1)\,U_q'\Big/\,\mathfrak{I}_q^{\,!} \,\in\, \big\{0,1\big\} \tag{5.5}$$

We can clearly reduce to the case when $\epsilon(\overline{a}) = 0$: in fact, if $\overline{a}^{\,2} = \overline{a}$ then $\epsilon(\overline{a})$ is necessarily 0 or 1 (for it is unipotent too), and in the latter case we then find that

$\overline{a}_0 := 1 - \overline{a}$ is idempotent and $\epsilon(\overline{a}_0) = 0$. Also the lift $a \in U_q' \Big/ \mathfrak{I}_q^!$ can be chosen, in this case, such that: $\epsilon(a) = 0$. To simplify notation, we set $H := U_q \Big/ \mathfrak{I}_q$ and $H' := U_q' \Big/ \mathfrak{I}_q^!$. We shall prove that, if $a \in H'$, $\epsilon(a) = 0$ and $a^2 \equiv a \mod (q-1)\, H'$, then $a \equiv 0 \mod (q-1)\, H'$, i.e. $a \in (q-1)\, H'$; in fact, this will give (5.5).

Having assumed that $\mathfrak{I}_q$ to be strict, H' identifies with a $\mathbb{C}\big[q, q^{-1}\big]$–submodule of H given in terms of the coalgebra structure of the latter: the embedding is the one canonically induced by the maps $U_q' \longhookrightarrow U_q \relbar\joinrel\twoheadrightarrow U_q \Big/ \mathfrak{I}_q$. In fact, the kernel of the latter map is $U_q' \bigcap \mathfrak{I}_q$ (by strictness assumption). It is easy to see from definitions that $U_q' \bigcap \mathfrak{I}_q = \mathfrak{I}_q^!$. Thus H' does embed into H:

$$H' = \left\{ \eta \in H \;\middle|\; \delta_n(\eta) \in (q-1)^n \, H^{\otimes n} \,,\ \forall\, n \in \mathbb{N} \right\}. \tag{5.6}$$

Now, $a^2 \equiv a \mod (q-1)H'$ means $a = a^2 + (q-1)\, c$ for some $c \in H'$; since $\epsilon(a) = 0$, we have $\epsilon(c) = 0$ as well. Applying δ_n to the identity $a = a^2 + (q-1)\, c$ and using formula (4.6) we get

$$\delta_n(a) \;=\; \delta_n\big(a^2\big) \;+\; (q-1)\, \delta_n(c) \;=\; \sum_{\Lambda \cup Y = \{1,\dots,n\}} \delta_\Lambda(a)\; \delta_Y(a) \;+\; (q-1)\; \delta_n(c)$$

for all $n \in \mathbb{N}$, which — noting that $\delta_0(a) := \epsilon(a) = 0$ yields:

$$\delta_n(a) \;=\; \sum_{\substack{\Lambda \cup Y = \{1,\dots,n\} \\ \Lambda, Y \neq \emptyset}} \delta_\Lambda(a)\; \delta_Y(a) \;+\; (q-1)\; \delta_n(c) \tag{5.7}$$

Since $c \in H'$, the last summand $(q-1)\, \delta_n(c)$ in right-hand side of (5.7) belongs to $(q-1)^{n+1}\, H^{\otimes n}$, thanks to (5.6). Similarly, since $a \in H'$ we have $\delta_k(a) \in (q-1)^k\, H^{\otimes k}$ for all $k \in \mathbb{N}$, by (5.6) again: therefore each summand $\delta_\Lambda(a)\, \delta_Y(a)$ in right-hand side of (5.7) belongs to $(q-1)^{n+1}\, H^{\otimes n}$ as well. But then (5.7) yields $\delta_n(a) \in (q-1)^{n+1} H^{\otimes n}$ for all $n \in \mathbb{N}$, which, again by (5.6), means exactly that $a \in (q-1)\, H'$. This ends the proof of the first claim.

(2) We will use similar arguments to show this claim: $F[L] = F\big[G^*\big] \Big/ \mathcal{I}(L)$ has no non-trivial idempotents. Since $\mathfrak{C}_q^{\urcorner} = \mathcal{C}_q(L)$ and $\mathcal{C}(L) = \mathcal{C}\big(\widehat{L}\,\big)$, we can assume $L = \widehat{L}$, i.e. L is observable. This implies $\mathcal{I}(L) = \Psi\big(\mathcal{C}(L)\big)$, which is clearly the specialization at $q = 1$ of $\Psi\big(\mathcal{C}(L)\big) = U_q'\, \mathfrak{C}_q^{\urcorner}$; therefore, $F[L] = F\big[G^*\big] \Big/ \mathcal{I}(L)$ is canonically isomorphic to the specialization at $q = 1$ of $U_q' \Big/ U_q'\, \mathfrak{C}_q^{\urcorner}$.

From now on, one can mimic step by step the proof of part *(1)*. The only detail to modify is that one must take $U_q\, \mathfrak{C}_q^{\,+} =: \Psi(\mathfrak{C}_q)$ in place of $\mathfrak{I}_q$, and $U_q' \left(\mathfrak{C}_q^{\urcorner}\right)^+ =: \Psi\big(\mathfrak{C}_q^{\urcorner}\big)$ in place of $\mathfrak{I}_q^!$. Letting $H := U_q \Big/ \Psi(\mathfrak{C}_q)$, and $H' := U_q' \Big/ \Psi\big(\mathfrak{C}_q^{\urcorner}\big)$, the thesis amounts to prove that

$$a \in H' \,, \quad a^2 \equiv a \mod (q-1)\, H' \Rightarrow a \equiv 0 \mod (q-1)\, H'$$

(In fact also $a \equiv 1 \mod (q-1)\, H'$ would be ok, but, arguing as before, we'll restrict to the case $\epsilon(a) = 0$).

As $\mathfrak{C}_q$ is strict, it is easy to see from definitions that $\mathfrak{C}_q^{\Lsh} = U_q' \bigcap \mathfrak{C}_q$, hence $\Psi\big(\mathfrak{C}_q^{\Lsh}\big) := U_q' \, \big(\mathfrak{C}_q^{\Lsh}\big)^+ = U_q'\big(U_q' \cap \mathfrak{C}_q\big)^+$: the latter is the kernel of the map $U_q' \hookrightarrow U_q \twoheadrightarrow U_q \Big/ U_q \, \mathfrak{C}_q^{\,+}$, so H' embeds as a $\mathbb{C}\big[q, q^{-1}\big]$–submodule of H, namely

$$H' = \left\{ \eta \in H \;\middle|\; \delta_n(\eta) \in (q-1)^n \, H^{\otimes n} \, , \; \forall \, n \in \mathbb{N} \right\} .$$

With this description at hand, computations are as in the proof of claim *(1).* ☐

Our next results are about the behaviour of quantum subgroups under composition of Drinfeld-like maps.

Proposition 5.10. *Let* $\mathcal{I}_q \, , \, \mathcal{C}_q \, , \, \mathfrak{I}_q \, , \, \mathfrak{C}_q$ *be weak quantizations of a subgroup* K *of* G . *Then:*

1. $\mathcal{I}_q \subseteq \big(\mathcal{I}_q^{\curlyvee}\big)^! \;\; , \quad \mathcal{C}_q \subseteq \big(\mathcal{C}_q^{\triangledown}\big)^{\Lsh}$;

2. $\mathfrak{C}_q \supseteq \big(\mathfrak{C}_q^{\Lsh}\big)^{\triangledown} \;\; , \quad \mathfrak{I}_q \supseteq \big(\mathfrak{I}_q^{\,!}\big)^{\curlyvee}$.

Proof. *(1)* By the very definitions, for any $n \in \mathbb{N}$ we have

$$\delta_n\big(\mathcal{I}_q\big) \;\subseteq\; {J_{F_q}}^{\otimes n} \bigcap \Big(\textstyle\sum_{s=0}^n {F_q}^{\otimes s} \otimes \mathcal{I}_q \otimes {F_q}^{\otimes (n-s-1)} \Big) = \textstyle\sum_{s=0}^n {J_{F_q}}^{\otimes s} \otimes \mathcal{I}_q \otimes {J_{F_q}}^{\otimes (n-s-1)} =$$
$$= \; (q-1)^n \cdot \textstyle\sum_{s=0}^n \big(F_q^{\vee}\big)^{\otimes s} \otimes \mathcal{I}_q^{\curlyvee} \otimes \big(F_q^{\vee}\big)^{\otimes (n-s-1)}$$

which means exactly $\mathcal{I}_q \subseteq \big(\mathcal{I}_q^{\curlyvee}\big)^!$. Similarly we can remark that:

$$\delta_n\big(\mathcal{C}_q\big) \;\subseteq\; {J_{F_q}}^{\otimes n} \bigcap \big({F_q}^{\otimes (n-1)} \otimes \mathcal{C}_q \big) \;=\; {J_{F_q}}^{\otimes (n-1)} \otimes \big(\mathcal{C}_q \bigcap J_{F_q} \big) \;=\; (q-1)^n \big(F_q^{\vee}\big)^{\otimes (n-1)} \otimes \mathcal{C}_q^{\triangledown}$$

which means $\mathcal{C}_q \subseteq \big(\mathcal{C}_q^{\triangledown}\big)^{\Lsh}$. Therefore claim *(1)* is proved.

(2) As $\big(\mathfrak{C}_q^{\Lsh}\big)^{\triangledown}$ is generated — as an algebra — by $(q-1)^{-1} \mathfrak{C}_q^{\Lsh} \bigcap J_{U_q'}$, it is enough to show that the latter space is contained in $\mathfrak{C}_q$. Let, then, $x' \in \mathfrak{C}_q^{\Lsh} \bigcap J_{U_q'}$. Surely $\delta_1\big(x'\big) \in (q-1) \, \mathfrak{C}_q$, hence $x' = \delta_1\big(x'\big) + \epsilon\big(x'\big) \in (q-1) \, \mathfrak{C}_q$. Therefore $(q-1)^{-1} x' \in \mathfrak{C}_q$, q.e.d. Similarly, $\big(\mathfrak{I}_q^{\,!}\big)^{\curlyvee}$ is the left ideal of U_q' generated by $(q-1)^{-1} \, \mathfrak{I}_q^{\,!} \bigcap J_{U_q'}$, thus — since $U_q' \subseteq U_q$ — we must only prove that $(q-1)^{-1} \, \mathfrak{I}_q^{\,!} \bigcap J_{U_q'}$ is contained in U_q. Again, if $y' \in \mathfrak{I}_q^{\,!} \bigcap J_{U_q'}$ then $y' = \delta_1\big(y'\big) + \epsilon\big(y'\big) \in (q-1) \, \mathfrak{I}_q$. Thus we get $(q-1)^{-1} y' \in \mathfrak{I}_q$, and *(2)* is proved. ☐

Remarks:

(a) By repeated applications of the previous proposition it is easily proved that:

$$\mathcal{I}_q^{\curlyvee} = \Big(\big(\mathcal{I}_q^{\curlyvee}\big)^! \Big)^{\curlyvee} \;\; , \quad \mathcal{C}_q^{\triangledown} = \Big(\big(\mathcal{C}_q^{\triangledown}\big)^{\Lsh} \Big)^{\triangledown} \;\; , \quad \mathfrak{C}_q^{\Lsh} = \Big(\big(\mathfrak{C}_q^{\Lsh}\big)^{\triangledown} \Big)^{\Lsh} \;\; , \quad \mathfrak{I}_q^{\,!} = \Big(\big(\mathfrak{I}_q^{\,!}\big)^{\curlyvee} \Big)^!$$

(b) Since we proved that Drinfeld-like maps always produce *proper* quantizations, and that proper quantizations specialize to *coisotropic* subgroups (cf. Proposition 3.5), the following holds:

1. if $\mathcal{I}_q = \big(\mathcal{I}_q^{\curlyvee}\big)^!$ then $\mathcal{I}_q$ is a proper quantization (of type $\mathcal{I}$) of a coisotropic subgroup of G;
2. if $\mathcal{C}_q = \big(\mathcal{C}_q^{\triangledown}\big)^{\urcorner}$ then $\mathcal{C}_q$ is a proper quantization (of type $\mathcal{C}$) of a coisotropic subgroup of G;
3. if $\mathfrak{I}_q = \big(\mathfrak{I}_q^{!}\big)^{\curlyvee}$ then $\mathfrak{I}_q$ is a proper quantization (of type $\mathfrak{I}$) of a coisotropic subgroup of G;
4. if $\mathfrak{C}_q = \big(\mathfrak{C}_q^{\urcorner}\big)^{\triangledown}$ then $\mathfrak{C}_q$ is a proper quantization (of type $\mathfrak{C}$) of a coisotropic subgroup of G.

(c) Since the whole construction is independent of the existence of real structures all the above claims hold true in the *real* framework as well.

Next result reads as a converse of the previous one, holding for Drinfeld maps applied to *strict* quantizations:

Theorem 5.11.

(a) if $\mathcal{I}_q$ is a strict quantization of a coisotropic subgroup of G then $\mathcal{I}_q = \big(\mathcal{I}_q^{\curlyvee}\big)^!$;

(b) if $\mathcal{C}_q$ is a strict quantization of a coisotropic subgroup of G then $\mathcal{C}_q = \big(\mathcal{C}_q^{\triangledown}\big)^{\urcorner}$;

(c) if $\mathfrak{I}_q$ is a strict quantization of a coisotropic subgroup of G then $\mathfrak{I}_q = \big(\mathfrak{I}_q^{!}\big)^{\curlyvee}$;

(d) if $\mathfrak{C}_q$ is a strict quantization of a coisotropic subgroup of G then $\mathfrak{C}_q = \big(\mathfrak{C}_q^{\urcorner}\big)^{\triangledown}$;

(e) The above claims hold true in the real framework as well.

Proof. (a) Let $\mathcal{I}_q$ be a strict quantization; by Proposition 5.10*(1)*, it is enough to prove $\mathcal{I}_q \supseteq \big(\mathcal{I}_q^{\curlyvee}\big)^!$. For this we apply the argument used in [10], Proposition 4.3, to prove that $F_q \supseteq \big(F_q^{\vee}\big)'$.

We denote by L the closed, coisotropic, connected subgroup of G^* such that $\mathcal{I}_q^{\curlyvee} = \mathfrak{I}_q(L)$, as in Proposition 5.1, and with $\mathfrak{l}$ its Lie algebra.

Let $y' \in \big(\mathcal{I}_q^{\curlyvee}\big)^!$. Then there is $n \in \mathbb{N}$ and $y^{\vee} \in \mathcal{I}_q^{\curlyvee} \setminus (q-1)\,\mathcal{I}_q^{\curlyvee}$ such that $y' = (q-1)^n y^{\vee}$. As we have seen strictness of $\mathcal{I}_q$ implies strictness of $\mathcal{I}_q^{\curlyvee}$ and therefore $y^{\vee} \notin (q-1)\,F_q^{\vee}$, and so for $\overline{y^{\vee}} := y^{\vee} \mod (q-1)\,F_q^{\vee}$ we have $\overline{y^{\vee}} \neq 0 \in F_q^{\vee}\Big|_{q=1} = U\big(\mathfrak{g}^*\big)$.

As $F_q^{\vee}$ is a quantization of $U\big(\mathfrak{g}^*\big)$, we can pick an ordered basis $\{b_\lambda\}_{\lambda\in\Lambda}$ of $\mathfrak{g}^*$, and a subset $\big\{x_\lambda^{\vee}\big\}_{\lambda\in\Lambda}$ of $(q-1)^{-1}J_{F_q}$ so that $x_\lambda^{\vee} \mod (q-1)\,F_q^{\vee} = b_\lambda$ for all $\lambda \in \Lambda$; therefore $x_\lambda^{\vee} = (q-1)^{-1}x_\lambda$ for some $x_\lambda \in J_{F_q}$, for all λ (like in the proof of [10] Proposition 4.3). In addition, we choose now the basis and its lift so that a subset $\{b_\theta\}_{\theta\in\Theta}$ (for some suitable $\Theta \subseteq \Lambda$) is a basis of $\mathfrak{l}$, and, correspondingly, $\big\{x_\theta^{\vee}\big\}_{\theta\in\Theta} \subseteq \mathcal{I}_q^{\curlyvee}$. Since $\overline{y^{\vee}} \neq 0 \in F_q^{\vee}\big|_{q=1} = U\big(\mathfrak{g}^*\big)$, by the Poincaré-Birkhoff-Witt theorem there is a non-zero polynomial $P\big(\{b_\theta\}_{\theta\in\Theta}\big)$ in the b_θ's such that $\overline{y^\vee} = P\big(\{b_\theta\}_{\theta\in\Theta}\big)$, hence

$$y^\vee - P\big(\big\{x_\theta^\vee\big\}_{\theta\in\Theta}\big) \in \mathcal{I}_q^{\curlyvee} \bigcap (q-1)\, F_q^\vee = (q-1)\, \mathcal{I}_q^{\curlyvee} \quad .$$

This implies $y^\vee = P\big(\big\{x_\theta^\vee\big\}_{\theta\in\Theta}\big) + (q-1)^\nu\, y_1^\vee$ for some $\nu \in \mathbb{N}_+$ where we have $y_1^\vee \in \mathcal{I}_q^{\curlyvee} \setminus (q-1)\,\mathcal{I}_q^{\curlyvee}$.

One can see, like in [7], Lemma 4.12, that the polynomial P has degree not greater than n. Thus $y' = (q-1)^n y^\vee = (q-1)^n P\big(\big\{x_\theta^\vee\big\}_{\theta\in\Theta}\big) + (q-1)^{n+\nu} y_1^\vee$, and

$$(q-1)^n P\big(\big\{x_\theta^\vee\big\}_{\theta\in\Theta}\big) = (q-1)^n P\big(\big\{(q-1)^{-1}\, x_\theta\big\}_{\theta\in\Theta}\big) \in \mathcal{I}_q$$

by a degree argument. But now, Proposition 5.10 gives $\mathcal{I}_q \subseteq \big(\mathcal{I}_q^{\curlyvee}\big)^!$. Then

$$y_1' := y' - (q-1)^n P\big(\big\{x_\mu^\vee\big\}_{\theta\in\Theta}\big) \in \big(\mathcal{I}_q^{\curlyvee}\big)^! \quad \text{and} \quad y_1' = (q-1)^{n+\nu} y_1^\vee = (q-1)^{n_1} y_1^\vee$$

where $n_1 := n+\nu > n$, and $y_1^\vee \in \mathcal{I}_q^{\curlyvee} \setminus (q-1)\,\mathcal{I}_q^{\curlyvee}$. We can then repeat the construction, with y_1' instead of y', n_1 instead of n, etc.: iterating, we find an increasing sequence of numbers $\big\{n_s\big\}_{s\in\mathbb{N}}$ (with $n_0 := n$) and a sequence of polynomials $\big\{P_s\big(\{X_\theta\}_{\theta\in\Theta}\big)\big\}_{s\in\mathbb{N}}$ (again $P_0 := P$) such that the degree of $P_s\big(\{X_\theta\}_{\theta\in\Theta}\big)$ is at most n_s, and the formal identity $y' = \sum_{s\in\mathbb{N}} (q-1)^{n_s} P_s\big(\big\{x_\theta^\vee\big\}_{\theta\in\Theta}\big)$ holds.

Now set $I_n := \sum_{k=1}^{n} (q-1)^{n-k}\, \mathcal{I}_q^{\,k}$ (for all $n \in \mathbb{N}$), and let $\widehat{\mathcal{I}}_q$ be the topological completion of $\mathcal{I}_q$ with respect to the filtration provided by the I_n's. Then, by construction, $(q-1)^{n_s} P_s\big(\big\{x_\theta^\vee\big\}_{\theta\in\Theta}\big) \in I_n$ for all $s\in\mathbb{N}$. This yields

$$\sum_{s\in\mathbb{N}} (q-1)^{n_s} P_s\big(\big\{x_\theta^\vee\big\}_{\theta\in\Theta}\big) \in \widehat{\mathcal{I}}_q \qquad \text{and} \qquad y' = \sum_{s\in\mathbb{N}} (q-1)^{n_s} P_s\big(\big\{x_\theta^\vee\big\}_{\theta\in\Theta}\big)$$

where the last is an identity in $\widehat{\mathcal{I}}_q$. Thus $y' \in \big(\mathcal{I}_q^{\curlyvee}\big)^! \bigcap \widehat{\mathcal{I}}_q$. Again with the same arguments as in [10], we see that $\mathcal{I}_q \bigcap (q-1)^\ell\, \widehat{\mathcal{I}}_q = (q-1)^\ell\, \mathcal{I}_q$ for any $\ell \in \mathbb{N}$. This together with $y' \in \big(\mathcal{I}_q^{\curlyvee}\big)^! \bigcap \widehat{\mathcal{I}}_q$ give $y' = (q-1)^{-m} \eta$ for some $m \in \mathbb{N}$ and $\eta \in \mathcal{I}_q$; thus

$$\eta = (q-1)^m\, y' \in \mathcal{I}_q \bigcap (q-1)^m\, \widehat{\mathcal{I}}_q = (q-1)^m\, \mathcal{I}_q\,,$$

whence $y' \in \mathcal{I}_q$, q.e.d.

(b) Assume that $\mathcal{C}_q$ is a strict quantization; by Proposition 5.10*(2)*, it is enough to prove $\mathcal{C}_q \supseteq \big(\mathcal{C}_q^{\triangledown}\big)^{\urcorner}$. To do that, we resume the argument used in [10], Proposition 4.3, to show that $F_q \supseteq \big(F_q^\vee\big)'$.

We denote by L the closed, coisotropic, connected subgroup of G^* such that $\mathcal{C}_q^{\triangledown} = \mathfrak{C}_q(L)$ and with $\mathfrak{l}$ its Lie algebra.

Let $c' \in \big(\mathcal{C}_q^{\triangledown}\big)^{\urcorner}$. Then there exist $n \in \mathbb{N}$ and $c^\vee \in \mathcal{C}_q^{\triangledown} \setminus (q-1)\,\mathcal{C}_q^{\triangledown}$ such that $c' = (q-1)^n c^\vee$. Note that strictness of $\mathcal{C}_q$ implies strictness of $\mathcal{C}_q^{\triangledown}$; hence $c^\vee \notin (q-1)\, F_q^\vee$, so that for $\overline{c^\vee} := c^\vee \mod (q-1)\, F_q^\vee$ we have $\overline{c^\vee} \neq 0 \in F_q^\vee\big|_{q=1} = U\big(\mathfrak{g}^*\big)$. Moreover, $\overline{c^\vee} \in \mathcal{C}_q^{\triangledown}\big|_{q=1} = \mathfrak{C}(L) = U(\mathfrak{l}) \subseteq U\big(\mathfrak{g}^*\big)$.

Since $F_q^\vee$ is a quantization of $U\big(\mathfrak{g}^*\big)$, we can fix an ordered basis $\{b_\lambda\}_{\lambda\in\Lambda}$ of $\mathfrak{g}^*$, and a subset $\big\{x_\lambda^\vee\big\}_{\lambda\in\Lambda}$ of $(q-1)^{-1} J_{F_q}$ such that $x_\lambda^\vee \mod (q-1)\, F_q^\vee = b_\lambda$ for all $\lambda \in \Lambda$; so $x_\lambda^\vee = (q-1)^{-1} x_\lambda$ for some $x_\lambda \in J_{F_q}$, for all λ (as in the proof of [10] Proposition 4.3). We can choose both the basis and its lift so that a subset $\{b_\mu\}_{\mu \in M}$ is a basis of $\mathfrak{l}$ (here $M \subseteq \Lambda$), and, correspondingly, $\big\{x_\mu^\vee\big\}_{\mu \in M} \subseteq (q-1)^{-1} J_{F_q} \bigcap \mathcal{C}_q^{\triangledown}$. Since $\overline{c^\vee} \neq 0 \in F_q^\vee\Big|_{q=1} = U\big(\mathfrak{g}^*\big)$, by the Poincaré-Birkhoff-Witt theorem there exists a non-zero polynomial $P\big(\{b_\mu\}_{\mu \in M}\big)$ in variables b_μ's such that $\overline{c^\vee} = P\big(\{b_\mu\}_{\mu \in M}\big)$, hence:

$$c^\vee - P\big(\big\{x_\mu^\vee\big\}_{\mu \in M}\big) \, \in \, \mathcal{C}_q^{\triangledown} \bigcap \, (q-1) \, F_q^\vee \, = \, (q-1) \, \mathcal{C}_q^{\triangledown} \, .$$

Therefore, $c^\vee = P\big(\big\{x_\mu^\vee\big\}_{\mu \in M}\big) + (q-1)^\nu \, c_1^\vee$ for some $\nu \in \mathbb{N}_+$ where we have $c_1^\vee \in \mathcal{C}_q^{\triangledown} \setminus (q-1) \, \mathcal{C}_q^{\triangledown}$.

Now, we can see — like in [7], Lemma 4.12 — that the degree of P is not greater than n. Then

$$c' = (q-1)^n c^\vee = (q-1)^n P\big(\big\{x_\mu^\vee\big\}_{\mu \in M}\big) + (q-1)^{n+\nu} c_1^\vee$$

with $(q-1)^n P\big(\big\{x_\mu^\vee\big\}_{\mu \in M}\big) = (q-1)^n P\big(\big\{(q-1)^{-1} \, x_\mu\big\}_{\mu \in M}\big) \in \mathcal{C}_q$ because P has degree bounded (from above) by n. As $\mathcal{C}_q \subseteq \big(\mathcal{C}_q^{\triangledown}\big)^{\Lsh}$, by Proposition 5.10, we get

$$c_1' := c' - (q-1)^n P\big(\big\{x_\mu^\vee\big\}_{\mu \in M}\big) \in \big(\mathcal{C}_q^{\triangledown}\big)^{\Lsh} \quad \text{and} \quad c_1' = (q-1)^{n+\nu} c_1^\vee = (q-1)^{n_1} c_1^\vee$$

with $n_1 := n + \nu > n$, and $c_1^\vee \in \mathcal{C}_q^{\triangledown} \setminus (q-1) \, \mathcal{C}_q^{\triangledown}$. We can repeat this construction with c_1' in place of c', n_1 in place of n, etc.. Iterating, we get an increasing sequence of numbers $\big\{n_s\big\}_{s \in \mathbb{N}}$ $(n_0 := n)$ and a sequence of polynomials $\big\{P_s\big(\{X_\mu\}_{\mu \in M}\big)\big\}_{s \in \mathbb{N}}$ $(P_0 := P)$ such that the degree of $P_s\big(\{X_\mu\}_{\mu \in M}\big)$ is at most n_s, and $c' = \sum_{s \in \mathbb{N}} (q-1)^{n_s} P_s\big(\big\{x_\mu^\vee\big\}_{\mu \in M}\big)$.

Consider

$$I_{\mathcal{C}_q} := Ker\Big(\mathcal{C}_q \overset{\epsilon}{\twoheadrightarrow} \mathbb{C}\big[q, q^{-1}\big] \overset{ev_1}{\twoheadrightarrow} \mathbb{C}\Big) = Ker\Big(\mathcal{C}_q \overset{ev_1}{\twoheadrightarrow} \mathcal{C}_q \big/ (q-1) \, \mathcal{C}_q \overset{\bar{\epsilon}}{\twoheadrightarrow} \mathbb{C}\Big)$$

By construction, $(q-1)^{n_s} P_s\big(\big\{x_\mu^\vee\big\}_{\mu \in M}\big) \in {I_{\mathcal{C}_q}}^{n_s}$ for all $s \in \mathbb{N}$; this means that $\sum_{s \in \mathbb{N}} (q-1)^{n_s} P_s\big(\big\{x_\mu^\vee\big\}_{\mu \in M}\big) \in \widehat{\mathcal{C}_q}$, the latter being the $I_{\mathcal{C}_q}$–adic completion of $\mathcal{C}_q$, and the formal expression $c' = \sum_{s \in \mathbb{N}} (q-1)^{n_s} P_s\big(\big\{x_\mu^\vee\big\}_{\mu \in M}\big)$ is an identity in $\widehat{\mathcal{C}_q}$: therefore $c' \in \big(\mathcal{C}_q^{\triangledown}\big)^{\Lsh} \bigcap \widehat{\mathcal{C}_q}$. Acting as in [10], again, we see that $\mathcal{C}_q \bigcap \, (q-1)^\ell \, \widehat{\mathcal{C}_q} = (q-1)^\ell \, \mathcal{C}_q$ for all $\ell \in \mathbb{N}$. Getting back to $c' \in \big(\mathcal{C}_q^{\triangledown}\big)^{\Lsh} \bigcap \widehat{\mathcal{C}_q}$, we have $c' = (q-1)^{-m} \kappa$ for some $m \in \mathbb{N}$ and $\kappa \in \mathcal{C}_q$; thus $\kappa = (q-1)^m \, c' \in \mathcal{C}_q \bigcap \, (q-1)^m \, \widehat{\mathcal{C}_q} = (q-1)^m \, \mathcal{C}_q$, whence $c' \in \mathcal{C}_q$, q.e.d.

(c) Let $\mathfrak{I}_q$ be a strict quantization: by Proposition 5.10*(2)* it is enough to prove $\mathfrak{I}_q \subseteq \big(\mathfrak{I}_q^!\big)^\curlyvee$; so given $y \in \mathfrak{I}_q$, we must prove that $y \in \big(\mathfrak{I}_q^!\big)^\curlyvee$. Recall that $\mathfrak{I}_q \subseteq U_q = \big(U_q'\big)^\vee$, the last identity following from Theorem 4.1. By construction,

$$\big(U_q'\big)^\vee \, = \, \textstyle\sum_{n \geq 0} (q-1)^{-n} \, I_{U_q'}^{\,n} \quad , \qquad I_{U_q'} \, := \, \big(U_q'\big)^+ + (q-1) \, U_q'$$

so for $y \in \mathfrak{I}_q \subseteq U_q = \big(U_q'\big)^\vee$ there exists $N \in \mathbb{N}$ such that

$$y_+ \, := \, (q-1)^N \, y \, \in \, I_{U_q'}^{\,N} \, \subseteq \, U_q' \tag{5.8}$$

Strictness of $\mathfrak{I}_q$, i.e. $\mathfrak{I}_q \bigcap (q-1)\, U_q = (q-1)\, \mathfrak{I}_q$, implies

$$\Big(\sum_{s=1}^{n} U_q^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes U_q^{\otimes (n-s)} \Big) \bigcap \big((q-1)^n \, U_q^{\otimes n} \big) = (q-1)^n \, \Big(\sum_{s=1}^{n} U_q^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes U_q^{\otimes (n-s)} \Big)$$

for all $n \in \mathbb{N}_+$; then, by the very definitions, the latter yields $\mathfrak{I}_q^{\,!} = \mathfrak{I}_q \bigcap U_q'$.
If in (5.8) $N = 1$, then $y_+ = y \in U_q'$, thus $y \in \mathfrak{I}_q \bigcap U_q' = \mathfrak{I}_q^{\,!}$, q.e.d. If, instead, $N > 1$, then formula (5.8), along with $\mathfrak{I}_q \,\dot{\trianglelefteq}\, U_q$, yields:

$$\delta_n(y_+) \in \Big((q-1)^N \cdot \sum_{s=1}^{n} U_q^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes U_q^{\otimes (n-s)} \Big) \bigcap \big((q-1)^n \, U_q^{\otimes n} \big) \, , \quad \forall\, n \in \mathbb{N}_+ \quad (5.9)$$

and since $\mathfrak{I}_q$ is strict, from (5.9) one gets

$$\delta_n(y_+) \in (q-1)^n \sum_{s=1}^{n} U_q^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes U_q^{\otimes (n-s)} \qquad \forall\, n \in \mathbb{N}$$

which means $y_+ \in \mathfrak{I}_q^{\,!}$. Eventually, we have found $y_+ \in \mathfrak{I}_q^{\,!} \bigcap I_{U_q'}^{\,N}$.

Now look at $I_{\mathfrak{I}_q^!} := I_{U_q'} \bigcap \mathfrak{I}_q^{\,!}$. Using the fact that $U_q' = U_q(\mathfrak{g})' = F\big[G^*\big]$ — from Theorem 4.1 — and $\mathfrak{I}_q^{\,!} = \mathfrak{I}_q(K)^! = \mathcal{I}_q(L)$ for some coisotropic subgroup L in G^* — as granted by Proposition 5.5 — and still taking into account strcitness, by an easy geometrical argument (via specialization at $q = 1$) we see that

$$I_{U_q'}^{\,n} \bigcap \mathfrak{I}_q^{\,!} \, \equiv \, I_{\mathfrak{I}_q^!}^{\,n} \quad \mathrm{mod}\ (q-1)\, U_q' \qquad \forall\, n \in \mathbb{N}_+ \, .$$

This, together with $\mathfrak{I}_q \bigcap (q-1)\, U_q = (q-1)\, \mathfrak{I}_q$, yields also

$$I_{U_q'}^{\,n} \bigcap \mathfrak{I}_q^{\,!} \, \equiv \, I_{\mathfrak{I}_q^!}^{\,n} \quad \mathrm{mod}\ (q-1)\, \mathfrak{I}_q^{\,!} \qquad \forall\, n \in \mathbb{N}_+$$

Finally, by suitable, iterated cancellation of factors $(q-1)$, which is possible because $\mathfrak{I}_q \bigcap (q-1) U_q = (q-1)\, \mathfrak{I}_q$, we eventually obtain

$$I_{U_q'}^{\,n} \bigcap \mathfrak{I}_q^{\,!} \, \equiv \, I_{\mathfrak{I}_q^!}^{\,n} \quad \mathrm{mod}\ (q-1)^n \, \mathfrak{I}_q^{\,!} \qquad \forall\, n \in \mathbb{N}_+ \, .$$

To sum up, we have $y_+ \in I_{U_q'}^{\,N} \bigcap \mathfrak{I}_q^{\,!} = I_{\mathfrak{I}_q^!}^{\,N}$; therefore, by definitions,

$$y = (q-1)^{-N} \, y_+ \, \in \, (q-1)^{-N} \, I_{\mathfrak{I}_q^!}^{\,N} \, \subseteq \, \big(\mathfrak{I}_q^{\,!} \big)^{\curlyvee} .$$

(d) Let $\mathfrak{C}_q$ be a strict quantization: by Proposition 5.10*(2)* it is enough to prove $\mathfrak{C}_q \subseteq \big(\mathfrak{C}_q^{\,\Lsh} \big)^{\triangledown}$. We follow the same arguments used for claim *(c)*. Let $c \in \mathfrak{C}_q$, since $\mathfrak{C}_q \subseteq U_q = \big(U_q' \big)^{\vee}$ — from Theorem 4.1 — and $\big(U_q' \big)^{\vee} = \sum_{n \geq 0} (q-1)^{-n} \, I_{U_q'}^{\,n}$, (notation as above) for $c \in \mathfrak{C}_q \subseteq U_q = \big(U_q' \big)^{\vee}$ there exists $N \in \mathbb{N}$ such that $c_+ := (q-1)^N \, c \, \in \, I_{U_q'}^{\,N} \, \subseteq \, U_q'$.

Now, strictness of $\mathfrak{C}_q$ implies

$$\big(U_q^{\otimes (n-1)} \otimes \mathfrak{C}_q \big) \bigcap (q-1)^n \, U_q^{\otimes n} = (q-1)^n \, \big(U_q^{\otimes (n-1)} \otimes \mathfrak{C}_q \big) \qquad \forall\, n \in \mathbb{N}_+$$

hence $\mathfrak{C}_q^{\,\Lsh} = \mathfrak{C}_q \bigcap U_q'$. If the above N is 1, then $c_+ = c \in U_q'$, thus $c \in \mathfrak{C}_q \bigcap U_q' = \mathfrak{C}_q^{\,\Lsh}$, q.e.d. If instead $N > 1$, then

$$\delta_n(c_+) \, \in \, \big((q-1)^N \cdot U_q^{\otimes n-1} \otimes \mathfrak{C}_q \big) \bigcap \big((q-1)^n \, U_q^{\otimes n} \big) \qquad \forall\, n \in \mathbb{N}_+$$

and, since $\mathfrak{C}_q$ is strict, $\delta_n(c_+) \in (q-1)^n \cdot U_q^{\otimes n-1} \otimes \mathfrak{C}_q$ for all $n \in \mathbb{N}_+$, which means $c_+ \in \mathfrak{C}_q^{\,\Lsh}$. Thus, eventually, we have $c_+ \in \mathfrak{C}_q^{\,\Lsh} \bigcap I_{U_q'}^{\,N}$.

Let us look, now, at $I_{\mathfrak{C}_q^{\Lsh}} := I_{U_q'} \bigcap \mathfrak{C}_q^{\,\Lsh}$. Again in force of strictness of $\mathfrak{C}_q$, a geometrical argument (at $q=1$) as before leads us to

$$I_{U_q'}^{\,n} \bigcap \mathfrak{C}_q^{\,\Lsh} \;\equiv\; I_{\mathfrak{C}_q^{\Lsh}}^{\,n} \quad \mathrm{mod}\ (q-1)^n\, \mathfrak{C}_q^{\,\Lsh} \quad , \qquad \forall\, n \in \mathbb{N}_+$$

from which we conclude that $c_+ \in I_{U_q'}^{\,N} \bigcap \mathfrak{C}_q^{\,\Lsh} = I_{\mathfrak{C}_q^{\Lsh}}^{\,N}$. Therefore, by the very definitions,

$$c \;=\; (q-1)^{-N}\, c_+ \;\in\; (q-1)^{-N}\, I_{\mathfrak{C}_q^{\Lsh}}^{\,N} \;\subseteq\; \big(\mathfrak{C}_q^{\,\Lsh}\big)^{\triangledown} \quad , \qquad \text{q.e.d.}$$

(e) This is a direct consequence of claims from *(a)* through *(d)*.

(f) Once again, this is true because the whole construction is independent of the existence of real structures. □

It is now time to clarify how the coisotropic subgroup L of G^* is linked to the coisotropic subgroup K of G. We will give this relation in the weak quantization case first, and show how it improves under stronger hypothesis.

Theorem 5.12. *Let K be a subgroup of G, and let $\mathcal{I}_q(K)$, $\mathcal{C}_q(K)$, $\mathfrak{I}_q(K)$ and $\mathfrak{C}_q(K)$ be weak quantizations as in Definition 3.6. Then (with notation of Proposition 2.2)*

(a) $\mathcal{I}_q(K)^{\curlyvee} = \mathfrak{I}_q\big(K^{\langle\perp\rangle}\big)$ *;*

(b) $\mathcal{C}_q(K)^{\triangledown} = \mathfrak{C}_q\big(K^{\langle\perp\rangle}\big)$ *;*

(c) *if* $\mathfrak{I}_q(K) = \big(\mathfrak{I}_q(K)^{!}\big)^{\curlyvee}$ *, then* $\mathfrak{I}_q(K)^{!} = \mathcal{I}_q\big(K^{\langle\perp\rangle}\big)$*; in particular, this holds if the quantization* $\mathfrak{I}_q(K)$ *is strict;*

(d) *if* $\mathfrak{C}_q = \big(\mathfrak{C}_q(K)^{\Lsh}\big)^{\triangledown}$*, then* $\mathfrak{C}_q(K)^{\Lsh} = \mathcal{C}_q\big(K^{\langle\perp\rangle}\big)$*; in particular, this holds if the quantization* $\mathfrak{C}_q(K)$ *is strict;*

(e) *claims (a–d) hold as well in the framework of real quantum subgroups.*

Proof. (a) By Proposition 5.1 we already have $\mathcal{I}_q(K)^{\curlyvee} = \mathfrak{I}_q(L)$ for some subgroup $L \subseteq G^*$. In order to show that $L = K^{\langle\perp\rangle}$, we will proceed much like in the proof of $F_q^{\vee} \big/ (q-1)\, F_q^{\vee} \cong U(\mathfrak{g}^*)$, as given in [10], Theorem 4.7.

Let us fix a subset $\{j_1, \dots, j_n\}$ of J adapted to K as in the proof of Proposition 5.1. Let $J^\vee := (q-1)^{-1} J \subset F_q^{\vee}$ and $j^{\,\vee} := (q-1)^{-1} j$ for all $j \in J$. From the discussion in that proof, we argue also that $\big\{ (q-1)^{-|\underline{e}|} j^{\,\underline{e}} \mod (q-1)\, F_q^{\vee} \;\big|\; \underline{e} \in \mathbb{N}^{\,n} \big\}$, where $j^{\,\underline{e}} = \prod_{s=1}^{n} j_s^{\underline{e}(i)}$, is a $\mathbb{C}$–basis of $F_1^{\vee}$, and $\big\{ j_1^{\vee}, \dots, j_n^{\vee} \big\}$ is a $\mathbb{C}$–basis of $\mathfrak{t} = J^\vee \mod (q-1)\, F_q^{\vee}$.

Now, $j_\mu\, j_\nu - j_\nu\, j_\mu \in (q{-}1)\, J$ (for $\mu, \nu \in \{1, \dots, n\}$) implies that:

$$j_\mu\, j_\nu - j_\nu\, j_\mu \;=\; (q-1) \textstyle\sum_{s=1}^{n} c_s\, j_s \;+\; (q-1)^2\, \gamma_1 \;+\; (q-1)\, \gamma_2$$

for some $c_s \in \mathbb{C}\big[q, q^{-1}\big]$, $\gamma_1 \in J$ and $\gamma_2 \in J^2$. Therefore

$$\big[j_\mu^{\vee}, j_\nu^{\vee} \big] := j_\mu^{\vee}\, j_\nu^{\vee} - j_\nu^{\vee}\, j_\mu^{\vee} = \textstyle\sum_{s=1}^{n} c_s\, j_s^{\vee} + \gamma_1 + (q{-}1)\, \gamma_2^{\vee} \equiv \textstyle\sum_{s=1}^{n} c_s\, j_s^{\vee} \quad \mathrm{mod}\ (q{-}1)\, F_q^{\vee}$$

(where we posed $\gamma_2^\vee := (q-1)^{-2}\gamma_2 \in (q-1)^{-2}\big(J^\vee\big)^2 \subseteq F_q^\vee$) thus the subspace $\mathfrak{t} := J^\vee \mod (q-1)\, F_q^\vee$ is a Lie subalgebra of $F_1^\vee$. But then it should be $F_1^\vee \cong U(\mathfrak{t})$ as Hopf algebras, by the above description of $F_1^\vee$ and PBW theorem.

Now for the second step. The specialization map $\pi^\vee \colon F_q^\vee \twoheadrightarrow F_1^\vee = U(\mathfrak{t})$ actually restricts to $\eta : J^\vee \twoheadrightarrow \mathfrak{t} = J^\vee \Big/ J^\vee \bigcap \big((q-1)\, F_q^\vee\big) = J^\vee \Big/ \big(J + J^\vee J\big)$, because $J^\vee \bigcap \big((q-1)\, F_q^\vee\big) = J^\vee \bigcap (q-1)^{-1} I_{F_q}^{\,2} = J + J^\vee J$. Also, multiplication by $(q-1)^{-1}$ yields a $\mathbb{C}\big[q,q^{-1}\big]$–module isomorphism $\mu : J \overset{\cong}{\hookrightarrow\!\!\!\!\!\rightarrow} J^\vee$. Let $\rho : \mathfrak{m}_e \twoheadrightarrow \mathfrak{m}_e \big/ {\mathfrak{m}_e}^2 = \mathfrak{g}^*$ be the natural projection map, and $\nu : \mathfrak{g}^* \hookrightarrow \mathfrak{m}_e$ a section of ρ. The specialization map $\pi : F_q \twoheadrightarrow F_1$ restricts to a map $\pi' \colon J \twoheadrightarrow J \big/ (J \bigcap (q-1)\, F_q) = \mathfrak{m}_e$. Let's fix a section $\gamma : \mathfrak{m}_e \hookrightarrow J$ of π' and consider the composition $\sigma := \eta \circ \mu \circ \gamma \circ \nu : \mathfrak{g}^* \longrightarrow \mathfrak{t}$: this is a well-defined Lie bialgebra morphism, independent of the choice of ν and γ.

In the proof of Proposition 5.1 we made a particular choice for the subset $\{j_1, \dots, j_n\}$. As a consequence, the above analysis to prove that $\sigma : \mathfrak{g}^* \cong \mathfrak{t}$ shows also that the left ideal $\mathcal{I}_1^{\curlyvee} := \mathcal{I}_q^{\curlyvee} \mod (q-1)\, F_q^\vee$ of $U(\mathfrak{t})$ is generated by

$$\eta\big(\mathcal{I}_q^{\curlyvee}\big) = (\eta \circ \mu)\big(\mathcal{I}_q\big) = (\sigma \circ \rho \circ \pi)\big(\mathcal{I}_q\big) = \sigma\big(\rho(\mathcal{I}\,)\big) = \sigma\big(\mathfrak{k}^\perp\big) \, .$$

So $\mathcal{I}_1^{\curlyvee} = U(\mathfrak{g}^*) \cdot \mathfrak{k}^\perp = U(\mathfrak{g}^*) \cdot \langle \mathfrak{k}^\perp \rangle = \mathfrak{I}\big(K^{\langle \perp \rangle}\big)$ — where we are identifying $\mathfrak{g}^*$ with its image via σ — which eventually means $\mathfrak{l} = \langle \mathfrak{k}^\perp \rangle$.

(b) By Proposition 5.3 we have $\mathcal{C}_q(K)^{\triangledown} = \mathfrak{C}_q(L)$ for some coisotropic subgroup L in G^*. We must prove that $L = K^{\langle \perp \rangle}$. Once again, we mimic the procedure of the proof of Proposition 5.3, and we fix a subset $\big\{j_1, \dots, j_n\big\}$ of J as in the proof of such Proposition. Then, tracking the analysis we did there to prove that $\sigma : \mathfrak{g}^* \cong \mathfrak{t}$, we see also that the unital subalgebra $\mathcal{C}_1^{\triangledown} := \mathcal{C}_q^{\triangledown} \mod (q–1)\, F_q^\vee$ of $U\big(\mathfrak{g}^*\big)$ is generated by $\eta\big(\mathcal{C}_q^{\triangledown}\big) = (\mu \circ \eta)\big(\mathcal{C}_q\big) = (\sigma \circ \rho \circ \pi)\big(\mathcal{C}_q\big) = \sigma\big(\rho(\mathcal{C}\,)\big) = \sigma\big(\mathfrak{k}^\perp\big)$. Thus $\mathcal{C}_1^{\triangledown}$ is the subalgebra of $U\big(\mathfrak{g}^*\big)$ generated by $\mathfrak{k}^\perp$, hence $\mathcal{C}_1^{\triangledown} = \big\langle \mathfrak{k}^\perp \big\rangle_{Alg} = U\big(\big\langle \mathfrak{k}^\perp \big\rangle_{Lie}\big) = U\big(\mathfrak{k}^{\langle \perp \rangle}\big) = \mathfrak{C}\big(K^{\langle \perp \rangle}\big)$, which means $\mathfrak{l} = \langle \mathfrak{k}^\perp \rangle$, q.e.d.

(c) Thanks to Proposition 5.5 we already know that $\mathfrak{I}_q(K)^! = \mathcal{I}_q(L)$ for some coisotropic subgroup L in G^*. Again, we must prove that $L = K^{\langle \perp \rangle}$. Note that we can assume K to be connected, as its relationship with $\mathfrak{I}_q(K)$ passes through $\mathfrak{k}$ alone; thus in the end we simply have to prove that $\mathfrak{l} := \mathit{Lie}\,(L) = \mathfrak{k}^{\langle \perp \rangle} = \mathfrak{k}^\perp$, taking into account that $\mathfrak{k}^{\langle \perp \rangle} = \mathfrak{k}^\perp$ because $\mathfrak{k}$ is coisotropic, by a remark following Proposition 5.10.

By assumption $\mathfrak{I}_q(K) = \big(\mathfrak{I}_q(K)^!\,\big)^{\curlyvee}$; this and *(a)* together give

$$\mathfrak{I}_q(K) \; = \; \big(\mathfrak{I}_q(K)^!\,\big)^{\curlyvee} = \mathcal{I}_q(L)^{\curlyvee} \; = \; \mathfrak{I}_q\big(L^{\langle \perp \rangle}\big) \; = \; \mathfrak{I}_q\big(L^\perp\big)$$

where $L^{\langle \perp \rangle} = L^\perp$ because L is coisotropic as well: at $q = 1$, this implies $\mathfrak{k} = \mathfrak{l}^\perp$, q.e.d.

(d) We must prove that $L = K^{\langle \perp \rangle}$: as above we can assume K to be connected, so we only have to prove that $\mathfrak{l} := \mathit{Lie}\,(L) = \mathfrak{k}^{\langle \perp \rangle} = \mathfrak{k}^\perp$ (as $\mathfrak{k}$ is coisotropic, by Proposition 5.11.

By assumption $\mathfrak{C}_q = \big(\mathfrak{C}_q(K)^{\urcorner}\big)^{\triangledown}$; this along with *(c)* gives

$$\mathfrak{C}_q(K) \; = \; \big(\mathfrak{C}_q(K)^{\urcorner}\big)^{\triangledown} = \mathcal{C}_q(L)^{\triangledown} \; = \; \mathfrak{C}_q\big(L^{\langle \perp \rangle}\big) \; = \; \mathfrak{C}_q\big(L^\perp\big)$$

with $L^{\langle\perp\rangle} = L^{\perp}$ since L is coisotropic too: specializing at $q=1$, this yields $\mathfrak{k} = \mathfrak{l}^{\perp}$.

(e) This is clear again since all arguments pass through unchanged in the real setup. □

Corollary 5.13. *Let $\mathcal{I}_q(K)$ and $\mathcal{C}_q(K)$ be weak quantizations of a (not necessarily) coisotropic subgroup K of G, of type $\mathcal{I}$ and $\mathcal{C}$ respectively. Then, with notation of Definition 2.1, we have*

$$\big(\mathcal{I}_q(K)^{\curlyvee}\big)^{!} = \mathcal{I}_q\big(\,\mathring{K}\,\big) \;\; , \qquad \big(\mathcal{C}_q(K)^{\triangledown}\big)^{\Lsh} = \mathcal{C}_q\big(\,\mathring{K}\,\big) \;\; .$$

Proof. Theorem 5.12*(a)* gives $\mathcal{I}_q(K)^{\curlyvee} = \mathfrak{I}_q\big(K^{\langle\perp\rangle}\big)$, and Proposition 5.10 yields

$$\Big(\mathfrak{I}_q\big(K^{\langle\perp\rangle}\big)^{!}\Big)^{\curlyvee} = \;\Big(\big(\mathcal{I}_q(K)^{\curlyvee}\big)^{!}\Big)^{\curlyvee} = \;\mathcal{I}_q(K)^{\curlyvee} \;=\; \mathfrak{I}_q\big(K^{\langle\perp\rangle}\big)$$

so that $\Big(\mathfrak{I}_q\big(K^{\langle\perp\rangle}\big)^{!}\Big)^{\curlyvee} = \;\mathfrak{I}_q\big(K^{\langle\perp\rangle}\big)$. Then Theorem 5.12 gives

$$\mathfrak{I}_q\big(K^{\langle\perp\rangle}\big)^{!} = \mathcal{I}_q\big((K^{\langle\perp\rangle})^{\langle\perp\rangle}\big) \;=\; \mathcal{I}_q\big(\mathring{\mathrm{K}}\,\big)$$

by Proposition 2.2. Therefore $\big(\mathcal{I}_q(K)^{\curlyvee}\big)^{!} = \mathfrak{I}_q\big(K^{\langle\perp\rangle}\big)^{!} = \mathcal{I}_q\big(\mathring{\mathrm{K}}\,\big)$ as claimed.

Similarly, Theorem 5.12*(b)* gives $\mathcal{C}_q(K)^{\triangledown} = \mathfrak{C}_q\big(K^{\langle\perp\rangle}\big)$, and the first remark after Proposition 5.10 yields

$$\Big(\mathfrak{C}_q\big(K^{\langle\perp\rangle}\big)^{\Lsh}\Big)^{\triangledown} = \;\Big(\big(\mathcal{C}_q(K)^{\triangledown}\big)^{\Lsh}\Big)^{\triangledown} = \;\mathcal{C}_q(K)^{\triangledown} \;=\; \mathfrak{C}_q\big(K^{\langle\perp\rangle}\big)$$

so that $\Big(\mathfrak{C}_q\big(K^{\langle\perp\rangle}\big)^{\Lsh}\Big)^{\triangledown} = \;\mathfrak{C}_q\big(K^{\langle\perp\rangle}\big)$. Then again by Theorem 5.12*(d)* we get

$$\mathfrak{C}_q\big(K^{\langle\perp\rangle}\big)^{\Lsh} = \mathcal{C}_q\big((K^{\langle\perp\rangle})^{\langle\perp\rangle}\big) \;=\; \mathcal{C}_q\big(\mathring{\mathrm{K}}\,\big)$$

still by Proposition 2.2. Thus $\big(\mathcal{C}_q(K)^{\triangledown}\big)^{\Lsh} = \mathfrak{C}_q\big(K^{\langle\perp\rangle}\big)^{\Lsh} = \mathcal{C}_q\big(\mathring{\mathrm{K}}\,\big)$ as claimed. □

One might guess that this Corollary holds true for weak quantizations of type $\mathfrak{I}$ and $\mathfrak{C}$ as well: actually, we do not know whether this is true or false.

We now consider the "compatibility" among different Drinfeld-like maps acting on quantizations of different types over a single pair *(subgroup, space)*. Indeed, we show that Drinfeld's functors preserve the subgroup-space correspondence — Proposition 5.14 — and the orthogonality correspondence — Proposition 5.16 — (if either occurs at the beginning) between different quantizations as mentioned.

Proposition 5.14. *Let K be a closed subgroup of G, and let Ψ and Φ be the map mentioned in §2.1. Then the following holds:*

(a) Let $\mathcal{C}_q$ and $\mathcal{I}_q$ be as in Section 3. If $\Psi(\mathcal{C}_q) = \mathcal{I}_q$, then $\Psi\big(\mathcal{C}_q^{\triangledown}\big) = \mathcal{I}_q^{\curlyvee}$.

(b) Let $\mathcal{I}_q$ and $\mathcal{C}_q$ be as in Section 3. If $\Phi(\mathcal{I}_q) = \mathcal{C}_q$, then $\Phi\big(\mathcal{I}_q^{\curlyvee}\big) = \mathcal{C}_q^{\triangledown}$.

(c) Let $\mathfrak{C}_q$ *and* $\mathfrak{I}_q$ *be as in Section 3. If* $\Psi(\mathfrak{C}_q) = \mathfrak{I}_q$ *, then* $\Psi\big({\mathfrak{C}_q}^{\urcorner}\big) \subseteq {\mathfrak{I}_q}^{!}$.

(d) Let $\mathfrak{I}_q$ *and* $\mathfrak{C}_q$ *be as in Section 3. If* $\Phi(\mathfrak{I}_q) = \mathfrak{C}_q$ *, then* $\Phi\big({\mathfrak{I}_q}^{!}\big) = {\mathfrak{C}_q}^{\urcorner}$.

Proof. Claims *(a)* and *(c)* both follow trivially from definitions.

As to claim *(b)*, let $\eta \in \mathcal{C}_q^+ = \Phi(\mathcal{I}_q)^+$, so that $\Delta(\eta) \in \eta \otimes 1 + F_q \otimes \mathcal{I}_q$. Then $\eta^\vee := (q-1)^{-1}\eta$ enjoys

$$\Delta\big(\eta^\vee\big) \in \eta^\vee \otimes 1 + F_q \otimes (q-1)^{-1}\,\mathcal{I}_q \subseteq \eta^\vee \otimes 1 + {F_q}^{\vee} \otimes {\mathcal{I}_q}^{\curlyvee}$$

whence $\eta^\vee \in \big({F_q}^{\vee}\big)^{co\mathcal{I}_q^{\curlyvee}} =: \Phi\big({\mathcal{I}_q}^{\curlyvee}\big)$. Since ${\mathcal{C}_q}^{\triangledown}$ is generated (as a subalgebra) by $(q-1)^{-1}\,\mathcal{C}_q^+$, we conclude that ${\mathcal{C}_q}^{\triangledown} \subseteq \Phi\big({\mathcal{I}_q}^{\curlyvee}\big)$.

Conversely, let $\varphi \in \Phi\big({\mathcal{I}_q}^{\curlyvee}\big)$. Then $\Delta(\varphi) \in \varphi \otimes 1 + {F_q}^{\vee} \otimes {\mathcal{I}_q}^{\curlyvee}$, and there exists $n \in \mathbb{N}$ such that $\varphi_+ := (q-1)^n \varphi \in \mathcal{I}_q$, so that $\Delta(\varphi_+) \in F_q \otimes \mathcal{I}_q + \mathcal{I}_q \otimes F_q$ (since $\mathcal{I}_q \,\dot{\trianglelefteq}\, F_q$). Then

$$\Delta(\varphi_+) \in \big(\varphi_+ \otimes 1 + (q-1)^n {F_q}^{\vee} \otimes {\mathcal{I}_q}^{\curlyvee}\big) \bigcap \big(F_q \otimes \mathcal{I}_q + \mathcal{I}_q \otimes F_q\big)$$

or equivalently

$$\Delta(\varphi_+) \,-\, \varphi_+ \otimes 1 \;\in \big((q-1)^n {F_q}^{\vee} \otimes {\mathcal{I}_q}^{\curlyvee}\big) \bigcap \big(F_q \otimes \mathcal{I}_q + \mathcal{I}_q \otimes F_q\big) \tag{5.10}$$

Now, the description of ${\mathcal{I}_q}^{\curlyvee}$ given in the proof of Proposition 5.1 implies that

$$\big((q-1)^n {F_q}^{\vee} \otimes {\mathcal{I}_q}^{\curlyvee}\big) \bigcap \big(F_q \otimes \mathcal{I}_q + \mathcal{I}_q \otimes F_q\big) \;=\; F_q \otimes \mathcal{I}_q$$

this together with (5.10) yields $\Delta(\varphi_+) \in \varphi_+ \otimes 1 + F_q \otimes \mathcal{I}_q$, hence $\varphi_+ \in {F_q}^{co\mathcal{I}_q} =: \Phi(\mathcal{I}_q) = \mathcal{C}_q$ and so $\varphi \in (q-1)^n\,\mathcal{C}_q \bigcap {F_q}^{\vee}$. On the other hand, the description of ${\mathcal{C}_q}^{\triangledown}$ in the proof of Proposition 5.3 implies that $(q-1)^{-n}\,\mathcal{C}_q \bigcap {F_q}^{\vee} \subseteq {\mathcal{C}_q}^{\triangledown}$, hence we get $\varphi \in {\mathcal{C}_q}^{\triangledown}$, q.e.d.

We finish with claim *(d)*. For the inclusion $\Phi\big({\mathfrak{I}_q}^{!}\big) \supseteq {\mathfrak{C}_q}^{\urcorner}$, let $\kappa \in {\mathfrak{C}_q}^{\urcorner}$. Since $\Phi\big({\mathfrak{I}_q}^{!}\big)$ contains the scalars, we may assume that $\kappa \in \mathit{Ker}\,(\epsilon)$, thus $\Delta(\kappa) = \kappa \otimes 1 + 1 \otimes \kappa + \delta_2(\kappa)$. By Proposition 5.7, we have ${\mathfrak{C}_q}^{\urcorner} \dot{\trianglelefteq}_\ell\, U_q'$; thus $\Delta(\kappa) - \kappa \otimes 1 = 1 \otimes \kappa + \delta_2(\kappa) \in U_q' \otimes {\mathfrak{C}_q}^{\urcorner}$, and more precisely

$$\Delta(\kappa) - \kappa \otimes 1 = 1 \otimes \kappa + \delta_2(\kappa) \in U_q' \otimes \big({\mathfrak{C}_q}^{\urcorner}\big)^+ \quad .$$

Since ${\mathfrak{C}_q}^{\urcorner} \subseteq \Psi\big({\mathfrak{C}_q}^{\urcorner}\big) \subseteq {\mathfrak{I}_q}^{!}$, by claim *(c)*, we get $\Delta(\kappa) - \kappa \otimes 1 \in U_q' \otimes {\mathfrak{I}_q}^{!}$, so $\kappa \in \big(U_q'\big)^{co\mathfrak{I}_q^!} =: \Phi\big({\mathfrak{I}_q}^{!}\big)$. Thus ${\mathfrak{C}_q}^{\urcorner} \subseteq \Phi\big({\mathfrak{I}_q}^{!}\big)$. For the converse inclusion, let $\eta \in \Phi\big({\mathfrak{I}_q}^{!}\big)$; again, we can assume $\eta \in \mathit{Ker}\,(\epsilon)$ too. As ${\mathfrak{I}_q}^{!} \subseteq \mathfrak{I}_q$, we get $\eta \in \Phi\big({\mathfrak{I}_q}^{!}\big) \subseteq \Phi\big(\mathfrak{I}_q\big) = \mathfrak{C}_q$. Then $\delta_n(\eta) \in {U_q}^{\otimes n} \otimes \mathfrak{C}_q$ for all $n \in \mathbb{N}_+$, so

$$\delta_n(\eta) \in (q-1)^n \left(\sum_{s=1}^{n-1} U_q^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes U_q^{\otimes (n-s)}\right) \bigcap \big(U_q^{\otimes (n-1)} \otimes \mathfrak{C}_q\big) \subseteq (q-1)^n\, U_q^{\otimes (n-1)} \otimes \mathfrak{C}_q$$

hence $\delta_n(\eta) \in (q-1)^n\, U_q^{\otimes (n-1)} \otimes \mathfrak{C}_q$ $(n \in \mathbb{N}_+)$ and $\eta \in \mathfrak{C}_q$, which means $\eta \in {\mathfrak{C}_q}^{\urcorner}$. $\square$

Remark 5.15. The inclusion $\Psi\big({\mathfrak{C}_q}^{\urcorner}\big) \subseteq {\mathfrak{I}_q}^{!}$ of Proposition 5.14*(c)* is *not* an identity in general — indeed, *counterexamples do exist.*

Finally, we look at what happens when our Drinfeld-like recipes are applied to a pair of quantizations associated to a same subgroup / homogeneous spaces with respect to some fixed double quantization (in the sense of Section 3). The result reads as follows:

Proposition 5.16. *Let* $\big(F_q[G]\,, U_q(\mathfrak{g})\big)$ *be a double quantization of* $(G, \mathfrak{g})$ *. Then:*

(a) Let $\mathcal{C}_q$ *and* $\mathfrak{I}_q$ *be weak quantizations and assume that* $\mathcal{C}_q = {\mathfrak{I}_q}^{\perp}$ *and* $\mathfrak{I}_q = {\mathcal{C}_q}^{\perp}$ *. Then* $\mathfrak{I}_q^{\,!} = \big(\mathcal{C}_q^{\,\triangledown}\big)^{\perp}$ *and* $\mathcal{C}_q^{\,\triangledown} \subseteq \big(\mathfrak{I}_q^{\,!}\big)^{\perp}$ *. If, in addition, either one of* $\mathcal{C}_q$ *or* $\mathfrak{I}_q$ *is strict, then also* $\mathcal{C}_q^{\,\triangledown} = \big(\mathfrak{I}_q^{\,!}\big)^{\perp}$ *.*

(b) Let $\mathfrak{C}_q$ *and* $\mathcal{I}_q$ *be weak quantizations and assume that* $\mathcal{I}_q = {\mathfrak{C}_q}^{\perp}$ *and* $\mathfrak{C}_q = {\mathcal{I}_q}^{\perp}$ *. Then* $\mathfrak{C}_q^{\,\Lsh} = \big(\mathcal{I}_q^{\,\curlyvee}\big)^{\perp}$ *and* $\mathcal{I}_q^{\,\curlyvee} \subseteq \big(\mathfrak{C}_q^{\,\Lsh}\big)^{\perp}$ *. If, in addition, either one of* $\mathfrak{C}_q$ *or* $\mathcal{I}_q$ *is strict, then also* $\mathcal{I}_q^{\,\curlyvee} = \big(\mathfrak{C}_q^{\,\Lsh}\big)^{\perp}$ *.*

Proof. Both in claim *(a)* and in claim *(b)* the orthogonality relations between $\mathfrak{C}_q$ and $\mathcal{I}_q$ and between $\mathcal{C}_q$ and $\mathfrak{I}_q$ are considered w.r.t. the pairing between $F_q[G]$ and $U_q(\mathfrak{g})$, and the subsequent orthogonality relations are meant w.r.t. the pairing between $F_q[G]^\vee$ and $U_q(\mathfrak{g})'$. Indeed, by Theorem 4.1, $\Big(U_q(\mathfrak{g})'\,, F_q[G]^\vee\Big)$ is a double quantization of $\big(G^*,\, \mathfrak{g}^*\big)$.

(a) First, $\epsilon(\mathfrak{I}_q) = 0$ because $\mathfrak{I}_q$ is a coideal. Then $x = \delta_1(x) \in (q-1)\,U_q$ for all $x \in \mathfrak{I}_q^{\,!}$, hence $\mathfrak{I}_q^{\,!} \subseteq (q-1)\,U_q$. Thus we have

$$\big\langle \mathcal{C}_q, \mathfrak{I}_q^{\,!} \big\rangle \subseteq (q-1)\,\mathbb{C}\big[q, q^{-1}\big] \quad .$$

Now let $J = J_{F_q}$ be the ideal of F_q, and take $c_i \in \mathcal{C}_q \cap J$ $(i = 1, \dots, n)$; then $\langle c_i, 1 \rangle = \epsilon(c_i) = 0$ $(i = 1, \dots, n)$. Given $y \in \mathfrak{I}_q^{\,!}$, look at

$$\Big\langle \prod_{i=1}^n c_i\,,\, y \Big\rangle = \Big\langle \bigotimes_{i=1}^n c_i\,, \Delta^n(y) \Big\rangle = \Big\langle \bigotimes_{i=1}^n c_i\,, \sum_{\Psi \subseteq \{1,\dots,n\}} \delta_\Psi(y) \Big\rangle = \sum_{\Psi \subseteq \{1,\dots,n\}} \Big\langle \bigotimes_{i=1}^n c_i\,, \delta_\Psi(y) \Big\rangle$$

Consider the summands in the last term of the above formula. Let $|\Psi| = t$ $(t \leq n)$, then

$$\big\langle \otimes_{i=1}^n c_i\,,\, \delta_\Psi(y) \big\rangle \;=\; \big\langle \otimes_{i \in \Psi} c_i\,,\, \delta_t(y) \big\rangle \cdot \prod_{j \notin \Psi} \langle c_j\,, 1 \rangle$$

by definition of δ_Ψ . Thanks to the previous analysis, we have $\prod_{j \notin \Psi} \langle c_j\,, 1 \rangle = 0$ unless $\Psi = \{1, \dots, n\}$, and in the latter case

$$\delta_\Psi(y) \;=\; \delta_n(y) \;\in\; (q-1)^n \sum_{s=1}^n {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)} \quad .$$

The outcome is

$$\Big\langle \prod_{i=1}^n c_i\,,\, y \Big\rangle = \Big\langle \bigotimes_{i=1}^n c_i\,,\, \delta_n(y) \Big\rangle \in \Big\langle \bigotimes_{i=1}^n c_i\,,\, (q-1)^n \sum_{s=1}^n {U_q}^{\otimes (s-1)} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes (n-s)} \Big\rangle = 0$$

because $y \in \mathfrak{I}_q^{\,!}$ and $\mathfrak{I}_q = {\mathcal{C}_q}^{\perp}$ by assumption. Thus $\big\langle (q-1)^{-n} \big(\mathcal{C}_q \bigcap J \big)^n , \, \mathfrak{I}_q^{\,!} \big\rangle = 0$, for all $n \in \mathbb{N}_+$. In addition, $\big\langle 1 \, , \, \mathfrak{I}_q^{\,!} \big\rangle = \epsilon\big(\mathfrak{I}_q^{\,!}\big) = 0$. The outcome is $\big\langle \mathcal{C}_q^{\triangledown} , \, \mathfrak{I}_q^{\,!} \big\rangle = 0$, whence $\mathfrak{I}_q^{\,!} \subseteq \big(\mathcal{C}_q^{\triangledown}\big)^{\perp}$, and $\mathcal{C}_q^{\triangledown} \subseteq \big(\mathfrak{I}_q^{\,!}\big)^{\perp}$.

Now we prove also $\big(\mathcal{C}_q^{\triangledown}\big)^{\perp} \subseteq \mathfrak{I}_q^{\,!}$. Notice that $\mathcal{C}_q^{\triangledown} \supseteq \mathcal{C}_q$, whence $\big(\mathcal{C}_q^{\triangledown}\big)^{\perp} \subseteq {\mathcal{C}_q}^{\perp} = \mathfrak{I}_q$; therefore $\big(\mathcal{C}_q^{\triangledown}\big)^{\perp} \subseteq \mathfrak{I}_q$. Pick now $\eta \in \big(\mathcal{C}_q^{\triangledown}\big)^{\perp}$ (inside ${U_q}'$). Since $\eta \in {U_q}'$, for all $n \in \mathbb{N}_+$ we have $\delta_n(\eta) \in (q-1)^n {U_q}^{\otimes n}$, and from $\eta \in \big(\mathcal{C}_q^{\triangledown}\big)^{\perp}$ we get also that $\eta_+ := (q-1)^{-n} \delta_n(\eta)$ enjoys $\Big\langle \big(\mathcal{C}_q \bigcap J_{F_q} \big)^{\otimes n} , \, \eta_+ \Big\rangle = 0$ — acting as before — so that

$$\eta_+ \in \Big(\big(\mathcal{C}_q \bigcap J_{F_q} \big)^{\otimes n} \Big)^{\perp} \; = \sum_{r+s=n-1} {U_q}^{\otimes r} \otimes \big(\mathcal{C}_q \bigcap J_{F_q} \big)^{\perp} \otimes {U_q}^{\otimes s} \quad .$$

Moreover $\delta_n(\eta) \in {J_{U_q}}^{\otimes n}$, hence $\delta_n(\eta) \in \big((q-1)^n {U_q}^{\otimes n} \big) \bigcap {J_{U_q}}^{\otimes n} = \; (q-1)^n {J_{U_q}}^{\otimes n}$, so

$$\begin{aligned} \eta_+ \in \Big(\big(\mathcal{C}_q \bigcap J_{F_q} \big)^{\otimes n} \Big)^{\perp} \bigcap {J_{U_q}}^{\otimes n} &= \Big(\sum_{r+s=n-1} {U_q}^{\otimes r} \otimes \big(\mathcal{C}_q \bigcap J_{F_q} \big)^{\perp} \otimes {U_q}^{\otimes s} \Big) \bigcap {J_{U_q}}^{\otimes n} = \\ &= \textstyle\sum_{r+s=n-1} {J_{U_q}}^{\otimes r} \otimes \Big(\big(\mathcal{C}_q \bigcap J_{F_q} \big)^{\perp} \bigcap J_{U_q} \Big) \otimes {J_{U_q}}^{\otimes s} \quad . \end{aligned}$$

Since $\big(\mathcal{C}_q \bigcap J_{F_q} \big)^{\perp} \bigcap \, J_{U_q} = {\mathcal{C}_q}^{\perp} \bigcap J_{U_q} = \mathfrak{I}_q \bigcap J_{U_q} = \mathfrak{I}_q$, we have

$$\eta_+ \; \in \sum_{r+s=n-1} {J_{U_q}}^{\otimes r} \otimes \mathfrak{I}_q \otimes {J_{U_q}}^{\otimes s}$$

whence

$$\delta_n(\eta) \; \in \; (q-1)^n \sum_{r+s=n-1} {U_q}^{\otimes r} \otimes \mathfrak{I}_q \otimes {U_q}^{\otimes s} \qquad \forall \; n \in \mathbb{N}_+ \quad .$$

Being, in addition, $\eta \in \mathfrak{I}_q$, for we proved that $\big(\mathcal{C}_q^{\triangledown}\big)^{\perp} \subseteq \mathfrak{I}_q$, we get $\eta \in \mathfrak{I}_q^{\,!}$. Therefore $\big(\mathcal{C}_q^{\triangledown}\big)^{\perp} \subseteq \mathfrak{I}_q^{\,!}$, q.e.d.

Finally, assume that $\mathcal{C}_q$ or $\mathfrak{I}_q$ are strict quantizations. Then we must still prove that $\mathcal{C}_q^{\triangledown} = \big(\mathfrak{I}_q^{\,!}\big)^{\perp}$. Since $\mathcal{C}_q = {\mathfrak{I}_q}^{\perp}$ and $\mathfrak{I}_q = {\mathcal{C}_q}^{\perp}$, it is easy to check that $\mathcal{C}_q$ is strict if and only if $\mathfrak{I}_q$ is; therefore, we can assume that $\mathfrak{I}_q$ is strict.

The assumptions and Theorem 5.11 *(b)* give $\mathfrak{I}_q = \big(\mathfrak{I}_q^{\,!}\big)^{\curlyvee}$; moreover, $\mathcal{I}_q := \mathfrak{I}_q^{\,!}$ is strict. Then we can apply the first part of claim *(b)* — which is proved, later on, in a way independent of the present proof of claim *(a)* itself — and get $\big(\mathcal{I}_q^{\curlyvee}\big)^{\perp} = \big({\mathcal{I}_q}^{\perp}\big)^{\Lsh}$. Therefore

$$\mathcal{C}_q^{\triangledown} \; = \; \big({\mathfrak{I}_q}^{\perp}\big)^{\triangledown} \; = \; \Big(\Big(\big(\mathfrak{I}_q^{\,!}\big)^{\curlyvee} \Big)^{\perp} \Big)^{\triangledown} \; = \; \Big(\big(\mathcal{I}_q^{\curlyvee}\big)^{\perp} \Big)^{\triangledown} \; = \; \Big(\big({\mathcal{I}_q}^{\perp}\big)^{\Lsh} \Big)^{\triangledown} \quad . \tag{5.11}$$

Now, it is straightforward to prove that $\mathcal{I}_q$ strict implies that ${\mathcal{I}_q}^{\perp}$ is strict as well. Then Proposition 5.11*(d)* ensures $\Big(\big({\mathcal{I}_q}^{\perp}\big)^{\Lsh} \Big)^{\triangledown} = {\mathcal{I}_q}^{\perp}$. This along with (5.11) yields $\mathcal{C}_q^{\triangledown} = \Big(\big({\mathcal{I}_q}^{\perp}\big)^{\Lsh} \Big)^{\triangledown} = {\mathcal{I}_q}^{\perp} = \big(\mathfrak{I}_q^{\,!}\big)^{\perp}$, ending the proof of *(a)*.

(b) With much the same arguments as for *(a)*, we find as well that

$$\big\langle \mathcal{I}_q^{\curlyvee} , \, \mathfrak{C}_q^{\Lsh} \big\rangle \; \in \; \big\langle J^{\otimes (n-1)} \otimes \mathcal{I}_q \, , \, {U_q}^{\otimes (n-1)} \otimes \mathfrak{C}_q \big\rangle \; \subseteq \; \big\langle \mathcal{I}_q \, , \mathfrak{C}_q \big\rangle \; = \; 0$$

because ${\mathcal{I}_q} = {\mathfrak{C}_q}^{\perp}$; this means that

$$ {\mathcal{I}_q}^{\curlyvee} \subseteq \big({\mathfrak{C}_q}^{\Lsh} \big)^{\perp} \quad , \qquad\qquad {\mathfrak{C}_q}^{\Lsh} \subseteq \big({\mathcal{I}_q}^{\curlyvee} \big)^{\perp} \quad . \tag{5.12} $$

Let now $\kappa \in \big({\mathcal{I}_q}^{\curlyvee} \big)^{\perp}_q \; \big(\subseteq {U_q}' \big)$. Since $\kappa \in {U_q}'$, we have $\delta_n(\kappa) \in (q-1)^n {U_q}^{\otimes n}$ for all $n \in \mathbb{N}$; moreover, from $\kappa \in \big({\mathcal{I}_q}^{\curlyvee} \big)^{\perp}$ it follows that $\kappa_+ := (q-1)^{-n} \delta_n(\kappa) \in {U_q}^{\otimes n}$ enjoys $\Big\langle J^{\otimes (n-1)} \otimes \mathcal{I}_q \, , \, \kappa_+ \Big\rangle = 0$, so that

$$ \kappa_+ \in \Big(J^{\otimes (n-1)} \otimes \mathcal{I}_q \Big)^{\perp} = \textstyle\sum_{r+s=n-2} {U_q}^{\otimes r} \otimes J^{\perp} \otimes {U_q}^{\otimes s} \otimes U_q \, + \, {U_q}^{\otimes (n-1)} \otimes {\mathcal{I}_q}^{\perp} \;\; . $$

In addition, $\delta_n(\kappa) \in {J_{U_q}}^{\otimes n}$, where $J_{U_q} := \mathit{Ker} \big(\epsilon : \, U_q \longrightarrow \mathbb{C}\big[q,q^{-1}\big] \big)$; therefore $\delta_n(\kappa) \in \big((q-1)^n {U_q}^{\otimes n} \big) \bigcap {J_{U_q}}^{\otimes n} = (q-1)^n {J_{U_q}}^{\otimes n}$, which together with the above formula yields

$$ \begin{aligned} \kappa_+ \, \in \, & \Big(J^{\otimes (n-1)} \otimes \mathcal{I}_q \Big)^{\perp} \bigcap {J_{U_q}}^{\otimes n} \, = \\ = & \, \Bigg(\sum_{r+s=n-2} {U_q}^{\otimes r} \otimes J^{\perp} \otimes {U_q}^{\otimes s} \otimes U_q \Bigg) \bigcap {J_{U_q}}^{\otimes n} \, + \, \Big({U_q}^{\otimes (n-1)} \otimes {\mathcal{I}_q}^{\perp} \Big) \bigcap {J_{U_q}}^{\otimes n} \, = \\ & \quad = \sum_{r+s=n-2} {J_{U_q}}^{\otimes r} \otimes \Big(J^{\perp} \bigcap J_{U_q} \Big) \otimes {J_{U_q}}^{\otimes s} \otimes J_{U_q} \, + \, {J_{U_q}}^{\otimes (n-1)} \otimes \Big({\mathcal{I}_q}^{\perp} \bigcap J_{U_q} \Big) \, = \\ & \qquad = {J_{U_q}}^{\otimes (n-1)} \otimes \Big({\mathcal{I}_q}^{\perp} \bigcap J_{U_q} \Big) \, = \, {J_{U_q}}^{\otimes (n-1)} \otimes \Big(\mathfrak{C}_q \bigcap J_{U_q} \Big) \, \subseteq \, {U_q}^{\otimes (n-1)} \otimes \mathfrak{C}_q \end{aligned} $$

where in the third equality we used the fact that $J^{\perp} \bigcap J_{U_q} = \{0\}$. So $\kappa_+ \in {U_q}^{\otimes (n-1)} \otimes \mathfrak{C}_q$, hence $\delta_n(\kappa) \in (q-1)^n {U_q}^{\otimes (n-1)} \otimes \mathfrak{C}_q$ for all $n \in \mathbb{N}_+$: thus $\kappa \in {\mathfrak{C}_q}^{\Lsh}$. Therefore $\big({\mathcal{I}_q}^{\curlyvee} \big)^{\perp} \subseteq {\mathfrak{C}_q}^{\Lsh}$, which together with the right-hand side inequality in (5.12) gives ${\mathfrak{C}_q}^{\Lsh} = \big({\mathcal{I}_q}^{\curlyvee} \big)^{\perp}$.

In the end, suppose also that one between $\mathfrak{C}_q$ and $\mathcal{I}_q$ is strict. As $\mathcal{I}_q = {\mathfrak{C}_q}^{\perp}$ and $\mathfrak{C}_q = {\mathcal{I}_q}^{\perp}$, one sees easily that $\mathcal{I}_q$ is strict if and only if $\mathfrak{C}_q$ is; then we can assume that $\mathfrak{C}_q$ is strict. We want to show that ${\mathcal{I}_q}^{\curlyvee} = \big({\mathfrak{C}_q}^{\Lsh} \big)^{\perp}$.

The assumptions and Theorem 5.11*(d)* give $\mathfrak{C}_q = \big({\mathfrak{C}_q}^{\Lsh} \big)^{\triangledown}$. Moreover, we have that $\mathcal{C}_q$ is strict by Proposition 5.3*(3)* and Proposition 5.7 *(3)*. Then we can apply the first part of claim *(a)*, thus getting $\big({\mathcal{C}_q}^{\triangledown} \big)^{\perp} = \big({\mathcal{C}_q}^{\perp} \big)^{!}$. Therefore

$$ {\mathcal{I}_q}^{\curlyvee} \, = \, \big({\mathfrak{C}_q}^{\perp} \big)^{\curlyvee} \, = \, \Big(\Big(\big({\mathfrak{C}_q}^{\Lsh} \big)^{\triangledown} \Big)^{\perp} \Big)^{\curlyvee} \, = \, \Big(\big({\mathcal{C}_q}^{\triangledown} \big)^{\perp} \Big)^{\curlyvee} \, = \, \Big(\big({\mathcal{C}_q}^{\perp} \big)^{!} \Big)^{\curlyvee} \tag{5.13} $$

Now, one proves easily that $\mathcal{C}_q$ strict implies ${\mathcal{C}_q}^{\perp}$ strict. Then Theorem 5.11*(c)* yields $\Big(\big({\mathcal{C}_q}^{\perp} \big)^{!} \Big)^{\curlyvee} = {\mathcal{C}_q}^{\perp}$. This and (5.13) give ${\mathcal{I}_q}^{\curlyvee} = \Big(\big({\mathcal{C}_q}^{\perp} \big)^{!} \Big)^{\curlyvee} = {\mathcal{C}_q}^{\perp} = \big({\mathfrak{C}_q}^{\Lsh} \big)^{\perp}$, which ends the proof of *(b)*. □

6 Examples

In this last section we will give some examples showing how our general constructions may be explicitely implemented. Some of the examples may look rather singular, but our aim here is mainly to draw the reader's attention on how even badly behaved cases can produce reasonable results. It has to be remarked that a wealth of new examples of coisotropic subgroups of Poisson groups have been recently produced ([20]), to which our recipes could be interestely applied.

N.B.: for the next examples — Subsections 6.2 and 6.3 — one can perform the explicit computations (that we just sketch) using definitions, formulas and notations as in [4], §6, and in [9], §7.

6.1 Quantization of Stokes matrices as an SL_n^*–space

As a first example, we mention the following. A well-known structure of Poisson group is defined on SL_n , typically known as the *standard* one; then one can consider its (connected) dual Poisson group SL_n^* , which in turn is a Poisson group as well. The set of *Stokes matrices* — i.e. upper triangular, unipotent matrices — of size n bears a natural structure of Poisson homogeneous space, and even Poisson quotient, for SL_n^* . In [4], Section 6, it was shown that one can find an explicit quantization, of formal type, of this Poisson quotient by a suitable application of the QDP procedure for formal quantizations developed in that paper.

Now, let us look at the explicit presentation of the formal quantization $U_\hbar(\mathfrak{sl}_n)$ considered in [*loc. cit.*]. One sees easily that this can be turned in a presentation of a *global* quantization (of $\mathfrak{sl}_n$ again), i.e. a QUEA $U_q(\mathfrak{sl}_n)$ in the sense of Section 3. Similarly, Drinfeld's QDP (for quantum groups) applied to $U_\hbar(\mathfrak{sl}_n)$ provides a formal quantization $F_\hbar[[SL_n^*]] := U_\hbar(\mathfrak{sl}_n)'$ of the function algebra over the formal group SL_n^* ; but then the analogous functor for the global version of QDP yields (cf. Theorem 4.1) a global quantization $F_q[SL_n^*] := U_q(\mathfrak{sl}_n)'$ of the function algebra over SL_n^* . In a nutshell, $F_q[SL_n^*]$ is nothing but (a suitable renormalization of) an obvious $\mathbb{C}\big[q,q^{-1}\big]$–integral form of $F_\hbar[[SL_n^*]]$.

Carrying further on this comparison, one can easily see that the whole analysis performed in [4] can be converted into a similar analysis for the global context, yielding parallel results; in particular, *one ends up with a global quantization — of type $\mathcal{C}$, in the sense of Section 3 — of the space of Stokes matrices.* More in detail, this quantization is a strict one, as such is the quantum subobject one starts with.

Since all this does not require more than a word by word translation, we refrain from filling in details.

6.2 A parametrized family of real coisotropic subgroups

Coisotropic subgroups may came in families, in some cases inside the same conjugacy class (which is responsible for different Poisson homogeneous bivectors on the same underlying manifold). An example in the real case was described in detail in [1]. The setting is the one of standard Poisson $SL_2(\mathbb{R})$, which contains a two parameter family of 1–dimensional coisotropic subgroups described, globally, by the right ideal and two-sided ideal

$$\mathcal{I}_{\mu,\nu} := \Big\{ a - d + 2\, q^{\frac{1}{2}} \mu b \, , \; q\, \nu b + c \Big\} \cdot F_q\big[SL_2(\mathbb{R})\big] \tag{6.1}$$

where a, b, c, d are the usual matrix elements generating $F_q\big[SL_2(\mathbb{R})\big]$, with $*$– structure in which they are all real (thus $q^* = q^{-1}$) and $\mu, \nu \in \mathbb{R}$. The corresponding family of coisotropic subgroups of classical $SL_2(\mathbb{R})$ may be described as

$$K_{\mu,\nu} := \left\{ \begin{pmatrix} d - 2\mu b & b \\ -\nu b & d \end{pmatrix} \,\middle|\, b, d \in \mathbb{R}\, , \; d^2 + \nu b^2 = 1 \right\}$$

(adapting our main text arguments to the case of *right* quantum coisotropic subgroups, this is quite trivial and we will do it without further comments). The corresponding $SL_2(\mathbb{R})$–quantum homogeneous spaces have local description given as follows: $\mathcal{C}_{\mu,\nu}$ is the subalgebra generated by

$$\begin{aligned} z_1 &= q^{-\frac{1}{2}}(ac + \nu bd) + 2\mu bc \, , \quad z_2 \,=\, c^2 + \nu d^2 + 2\mu q^{-\frac{1}{2}} cd \, , \\ z_3 &= a^2 + \nu b^2 + 2\mu q^{-\frac{1}{2}} ab \, . \end{aligned} \tag{6.2}$$

Using commutation relations — see (12) in [2] — it is easily seen that $\mathcal{C}_{\mu,\nu}$ has a linear basis given by $\big\{ z_1^p z_2^q \, , \, z_1^p z_3^r \,\big|\, p, q, r \in \mathbb{N} \big\}$.

Proposition 6.1. *The subalgebra $\mathcal{C}_{\mu,\nu}$ is a right coideal of $F_q\big[SL_2(\mathbb{R})\big]$ and is a strict quantization — of type $\mathcal{C}$ — of $K_{\mu,\nu}$.*

Proof. The first statement is proven in [2]. As for the second we will first show that $z_1^p z_2^q \, , z_1^p z_3^r \notin (q-1) F_q\big[SL_2(\mathbb{R})\big]$ for any $p, q, r \in \mathbb{N}$. This may done by considering their expression in terms of the usual basis $\big\{ a^p b^r c^s \, , \, b^h c^k d^i \big\}$ of $F_q\big[SL_2(\mathbb{R})\big]$. In fact we do not need a full expression of monomials $z_1^p z_2^r$ or $z_1^p z_3^r$ in terms of this basis, which would lead to quite heavy computations. It is enough to remark that, for example, since

$$z_1^p z_2^r \;=\; \Big(q^{-\frac{1}{2}} ac + b(\nu d + 2\mu c) \Big)^p \Big(c^2 + (\nu d + 2\mu q^{-\frac{1}{2}} c) d \Big)^r$$

we can get an element multiple of $a^p c^{p+2r}$ only from $(ac) \cdot \cdots \cdot (ac) \cdot c \cdot \cdots \cdot c$, which is of the form $q^h a^p c^{p+2r} \notin F_q\big[SL_2(\mathbb{R})\big]$. Since no other elements may add up with this one, we have $z_1^p z_2^r \notin (q-1) F_q\big[SL_2(\mathbb{R})\big]$. A similar argument works for $z_1^p z_3^r$.

In a similar way we prove that any $\mathbb{C}\big[q, q^{-1}\big]$–linear combination of the $z_1^p z_2^q$'s and the $z_1^s z_3^r$'s is in $(q-1) F_q\big[SL_2(\mathbb{R})\big]$ if and only if all coefficients are in $(q-1)\mathbb{C}\big[q, q^{-1}\big]$. Therefore $\mathcal{C}_q$ is strict, q.e.d. $\square$

It makes therefore sense to compute $\mathcal{C}_{\mu,\nu}^{\,\triangledown}$; to this end, we can resume a detailed description of $U_q(\mathfrak{sl}_2^*) := F_q\big[SL_2(\mathbb{R})\big]^\vee$ — apart for the real structure, which is not really relevant here — from [9], §7.7. From our PBW-type basis we have that $\mathcal{C}_{\mu,\nu}^{\,\triangledown}$ is the subalgebra of $F_q\big[SL_2(\mathbb{R})\big]^\vee$ generated by the elements $\zeta_i := \frac{1}{q-1}\big(z_i - \varepsilon(z_i)\big) \in F_q\big[SL_2(\mathbb{R})\big]^\vee$ $(i = 1, 2, 3)$. Since we know that

$$H_+ := \frac{a-1}{q-1} \, , \qquad E := \frac{b}{q-1} \, , \qquad F := \frac{c}{q-1} \, , \qquad H_- := \frac{d-1}{q-1}$$

are algebra generators of $U_q(\mathfrak{sl}_2^*) := F_q\big[SL_2(\mathbb{R})\big]^\vee$, we deduce that

$$\begin{aligned} \frac{\zeta_1}{q-1} &= q^{-\frac{1}{2}}(F+\nu E) + (q-1)\left(q^{-\frac{1}{2}}H_+F + q^{-\frac{1}{2}}\nu E H_- + 2\mu EF\right) \\ \frac{\zeta_2-\nu}{q-1} &= 2\,(\nu H_- + \mu q^{-\frac{1}{2}}F) + (q-1)\left(F^2 + \nu H_-^2 + 2\mu q^{-\frac{1}{2}}FH_-\right) \\ \frac{\zeta_3-1}{q-1} &= 2\,(H_+ + \mu q^{-\frac{1}{2}}E) + (q-1)\left(H_+^2 + \nu E^2 + 2\mu q^{-\frac{1}{2}}H_+E\right) \end{aligned} \tag{6.3}$$

In the semiclassical specialization $U_q(\mathfrak{sl}_2^*) \xrightarrow{q\longrightarrow} U_q(\mathfrak{sl}_2^*)\Big/(q-1)U_q(\mathfrak{sl}_2^*)$ one has that $E \mapsto \mathrm{e}$, $F \mapsto \mathrm{f}$, $H_\pm \mapsto \pm\mathrm{h}$, where $\mathrm{h}, \mathrm{e}, \mathrm{f}$ are Lie algebra generators of $\mathfrak{sl}_2^*$; therefore the semiclassical limit of the right hand side of (6.3) is the Lie subalgebra generated by $\mathrm{f} + \nu\,\mathrm{e}$, $-\nu\,\mathrm{h} + \mu\,\mathrm{e}$, $\mathrm{h} + \mu\,\mathrm{e}$, or, equivalently, the 2–dimensional Lie subalgebra generated by $\mathrm{f} + \nu\,\mathrm{e}$ and $\mathrm{h} + \mu\,\mathrm{e}$ (the three elements above being linearly dependent) with relation $[\,\mathrm{h} + \mu\,\mathrm{e}, \mathrm{f} + \nu\,\mathrm{e}\,] = \mathrm{f} + \nu\,\mathrm{e}$. The quantization of this coisotropic subalgebra of $\mathfrak{sl}_2^*$ is therefore the subalgebra generated inside $U_q(\mathfrak{sl}_2^*)$ by the quadratic elements (6.3).

Similar computations can be performed starting from $\mathcal{I}_{\mu,\nu}$. The transformed $\mathcal{I}_{\mu,\nu}^{\curlyvee}$ is the right ideal generated by the image of $a - d + 2\,q^{\frac{1}{2}}\mu b$ and $q\nu b + c$, i.e. the right ideal generated by $H_+ - H_- + 2\,q^{\frac{1}{2}}\mu E$ and $q\,\nu E + F$; moreover, from its semiclassical limit, it is easily seen that this again corresponds to the same coisotropic subgroup of the dual Poisson group $SL_2(\mathbb{R})^*$.

All this gives a local — i.e., infinitesimal — description of the (2–dimensional) coisotropic subgroups $K_{\mu,\nu}^\perp$ in $SL_2(\mathbb{R})^*$.

6.3 The non coisotropic case

Let us finally consider the case of a non coisotropic subgroup. We will consider the embedding of $SL_2(\mathbb{C})$ into $SL_3(\mathbb{C})$ corresponding to a non simple root, which easily generalizes to higher dimensions. Computations will only be sketched.

Let $\mathfrak{h}$ be the subalgebra of $\mathfrak{sl}_3(\mathbb{C})$ spanned by $E_{1,3}$, $F_{1,3}$, $H_{1,3} = H_1 + H_2$. Easy computations show that the standard cobracket values are

$$\begin{aligned} \delta(E_{13}) &= E_{13} \wedge (H_1 + H_2) + 2E_{23} \wedge E_{12} \\ \delta(F_{13}) &= F_{13} \wedge (H_1 + H_2) - 2F_{23} \wedge F_{12} \\ \delta(H_1 + H_2) &= 0 \end{aligned} \tag{6.4}$$

and, therefore, the corresponding embedding $SL_2(\mathbb{C}) \hookrightarrow SL_3(\mathbb{C})$ is *not* coisotropic. To compute the coisotropic interior $\overset{\circ}{\mathfrak{h}}$ of $\mathfrak{h}$, consider that $\langle H_1 + H_2 \rangle$ is, trivially, a subbialgebra of $\mathfrak{h}$, thus contained in $\overset{\circ}{\mathfrak{h}}$. Let $X := (H_1 + H_2) + \alpha E_{13} + \beta F_{13}$: then

$$\delta(X) \;=\; X \wedge (H_1 + H_2) + 2\,(\alpha E_{23} \wedge E_{12} - \beta F_{23} \wedge F_{12})$$

shows that no such X is in $\overset{\circ}{\mathfrak{h}}$, unless $\alpha = 0 = \beta$. The oucome is that we have

$$\overset{\circ}{H} \;=\; \begin{pmatrix} \gamma & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \gamma^{-1} \end{pmatrix} \subseteq SL_3(\mathbb{C})$$

with $\gamma \in \mathbb{C}^*$. Correspondingly

$$\mathfrak{h}^{\langle\perp\rangle} \; = \; \Big(\overset{\circ}{\mathfrak{h}} \Big)^{\perp} \; = \; \big\langle\, \mathrm{e}_{1,2} \, , \, \mathrm{e}_{1,3} \, , \, \mathrm{e}_{2,3} \, , \, \mathrm{f}_{1,2} \, , \, \mathrm{f}_{1,3} \, , \, \mathrm{f}_{2,3} \, , \, \mathrm{h}_{2.2} \,\big\rangle \qquad \Big(\subseteq \mathfrak{sl}_3(\mathbb{C})^* \Big)$$

and, thus $SL_3(\mathbb{C})^* \Big/ H^{\langle\perp\rangle}$ is a 1–dimensional Poisson homogeneous spaces — with, of course, zero Poisson bracket.

Let us consider now any weak quantization $\mathfrak{C}_q(H)$ of H. It should certainly contain the subalgebra of $U_q(\mathfrak{sl}_3)$ generated by the root vectors $E_{1,3}$, $F_{1,3}$, together with $K_1 K_3^{-1}$ and $\widehat{H}_{1,3} := \big(K_1 K_3^{-1} - 1 \big) \Big/ (q-1)$. The equality

$$\Delta(E_{1,3}) \; = \; E_{1,3} \otimes K_1 K_3^{-1} + 1 \otimes E_{1,3} + (q-1) E_{1,2} \otimes E_{2,3}$$

tells us that, in order to be a left coideal, such a quantization should also contain either $(q-1)E_{1,2}$ or $(q-1)E_{2,3}$ (and thus, as expected, it cannot be strict). Let us try to compute some elements in $\mathfrak{C}_q(H)^{\Lsh}$. Certainly, since

$$\delta_2\big(\widehat{H}_{1,3}\big) \; = \; \widehat{H}_{1,3} \otimes \big(K_1 K_3^{-1} - 1 \big) \; = \; (q-1)\, \widehat{H}_{1,3} \otimes \widehat{H}_{1,3}$$

we can conclude that $(q-1)\widehat{H}_{1,3} \in \mathfrak{C}_q(H)^{\Lsh}$. On the other hand,

$$\delta_2(E_{1,3}) \; = \; (q-1) E_{1,3} \otimes \widehat{H}_{1,3} + (q-1) E_{1,2} \otimes E_{2,3}$$

implies that $(q-1)E_{1,3} \notin \mathfrak{C}_q(H)^{\Lsh}$, while $(q-1)^2 E_{1,3} \in \mathfrak{C}_q(H)^{\Lsh}$.

All this means the following.

Within $\mathfrak{C}_q(H)^{\Lsh}$ we find a non-diagonal matrix element of the form $(q-1)\, t_{1,3}$: it belong to $(q-1) U_q(\mathfrak{sl}_3)'$ but not to $(q-1)\mathfrak{C}_q(H)^{\Lsh}$, so that

$$\mathfrak{C}_q(H)^{\Lsh} \bigcap\, (q-1)\, U_q(\mathfrak{sl}_3)' \;\; \underset{\neq}{\supseteq} \;\; (q-1)\, \mathfrak{C}_q(H)^{\Lsh}$$

which means that the quantization $\mathfrak{C}_q(H)^{\Lsh}$ *is not strict.* On the other hand, we know by Proposition 5.7*(3)* that $\mathfrak{C}_q(H)^{\Lsh}$ is *proper*. Therefore, *we have an example of a quantization (of type $\mathcal{C}_q$, still by Proposition 5.7(3)) which is proper yet not strict.*

In addition, in the specialization map $\pi : U_q(\mathfrak{sl}_3)' \relbar\joinrel\twoheadrightarrow U_q(\mathfrak{sl}_3)' \Big/ (q-1) U_q(\mathfrak{sl}_3)'$ the element $(q-1)\, t_{1,3}$ is obviously mapped to zero, i.e. it yields a trivial contribution to the semiclassical limit of $\mathfrak{C}_q(H)^{\Lsh}$ — which here is meant as being $\pi\big(\mathfrak{C}_q(H)^{\Lsh}\big) = \mathfrak{C}_q(H)^{\Lsh} \Big/ \mathfrak{C}_q(H)^{\Lsh} \bigcap (q-1) U_q(\mathfrak{sl}_3)'$. With similar computations it is possible to prove, in fact, that the only generating element in $\mathfrak{C}(H)^{\Lsh}$ having a non-trivial semiclassical limit is $(q-1)\widehat{H}_{1,3}$. Therefore, through specialization at $q = 1$, from $\mathfrak{C}(H)^{\Lsh}$ one gets only $\pi\big(\mathfrak{C}_q(H)^{\Lsh}\big) = \mathbb{C}\big[t_{2,2}\big]$: indeed, this in turn tells us exactly that $\mathfrak{C}_q(H)^{\Lsh}$ is a quantization, of *proper* type, of the homogeneous $SL_3(\mathbb{C})^*$–space $SL_3(\mathbb{C})^* \Big/ H^{\langle\perp\rangle}$ (whose Poisson bracket is trivial).

References

[1] F. Bonechi, N. Ciccoli, R. Giachetti, E. Sorace, M. Tarlini, *The coisotropic subgroup structure of $SL_q(2;\mathbb{R})$*, Journ. Geom. Phys. **37** (2001), 190–200.

[2] F. Bonechi, N. Ciccoli, M. Tarlini, *Noncommutative instantons on the 4–sphere from quantum groups*, Comm. Math. Phys. **226** (2002), 419–432.

[3] V. Chari, A. Pressley, *Quantum Groups*, Cambridge University Press, Cambridge (UK), 1994.

[4] N. Ciccoli, F. Gavarini, *A quantum duality principle for coisotropic subgroups and Poisson quotients*, Adv. Math. **199** (2006), 104–135.

[5] N. Ciccoli, R. Fioresi, F. Gavarini, *Quantization of projective homogeneous spaces and duality principle*, Journ. Noncomm. Geom. **2** (2008), 449–496.

[6] V. G. Drinfel'd, *Quantum groups*, Proc. Int. Congr. Math., Vol. 1, 2 (Berkeley, Calif., 1986), 798820, Amer. Math. Soc., Providence, RI, 1987.

[7] P. Etingof, V. Kazhdan, *Quantization of Lie bialgebras I*, Selecta Math. (N. S.) **2** (1996), 1–11.

[8] F. Gavarini, *The quantum duality principle*, Annales de l'Institut Fourier **52** (2002), no. 3, 809–834.

[9] F. Gavarini, *The global quantum duality principle: theory, examples, and applications*, preprint `arXiv:math/0303019v8 [math.QA]` (2003), 120 pages.

[10] F. Gavarini, *The global quantum duality principle*, Journ. für die Reine Angew. Math. **612** (2007), 17–33.

[11] F. Grosshans, *Algebraic homogeneous spaces and invariant theory*, Lect. Notes Math. **1673**, Springer, Berlin, 1997.

[12] I. Heckenberger, S. Kolb, *Right coideal subalgebras of the Borel part of a quantized universal enveloping algebra*, Int. Math. Res. Not. IMRN 2011, no. 2, 419–451.

[13] V. K. Karchenko, A. V. Lara Sagahón, *Right coideal subalgebras in $U_q(sl_{n+1})$*, J. Algebra **319** (2008), 2571–2625.

[14] C. Kassel, V. Turaev, *Biquantization of Lie bialgebras*, Pac. Journ. Math. **195** (2000), 297–369.

[15] G. Letzter, *Coideal subalgebras and quantum symmetric pairs*, in: S. Montgomery, H.–J. Schneider (eds.) New Directions in Hopf Algebras MSRI Publ. **43** (2002), 117–165.

[16] H.–J. Lu, *Moment maps at the quantum level*, Comm. Math. Phys. **157** (1993) 389–404.

[17] S. Montgomery, *Hopf Algebras and their actions on rings*, CBMS Regional Conf. Series in Math **82**, Amer. Math. Soc., Providence (RI), 1993.

[18] H.–J. Schneider, *Principal homogeneous spaces for arbitrary Hopf algebras*, Israel J. Math. **72** (1990), 167–195.

[19] M. Takeuchi, *Quotient spaces for Hopf algebras*, Comm. Alg. **22** (1994), 2503–2523.

[20] M. Zambon, *A construction for coisotropic subalgebras of Lie bialgebras*, Journ. Pure Appl. Alg. **215**, 411–419 (2011).

[21] S. Zakrzewszki, *Poisson homogeneous spaces*, in: J. Lukierski, Z. Popowicz, J. Sobczyk (eds.), Quantum Groups (Karpacz 1994), PWN Warszaw, 629–639, 1995.